

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification</b> <sup>6</sup> : <b>C07D 251/66, 251/70, A61K 31/53, C07D 403/12, 405/12, 409/12</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/36410</b> <b>(43) International Publication Date:</b> 22 July 1999 (22.07.99)
<b>(21) International Application Number:</b> PCT/US99/00945 <b>(22) International Filing Date:</b> 13 January 1999 (13.01.99)  <b>(30) Priority Data:</b> 09/006,430 13 January 1998 (13.01.98) US  <b>(71) Applicant:</b> SCRIPTGEN PHARMACEUTICALS, INC. [US/US]; 610 Lincoln Street, Waltham, MA 02154 (US).  <b>(72) Inventors:</b> ARENAS, Jaime, E.; 26 Ledgeawn Avenue, Lexington, MA 02173 (US). CLOAD, Sharon, T.; 162 Chilton Street, Cambridge, MA 02138 (US). FLEMING, Elizabeth, S.; 30 Shean Road, Belmont, MA 02178 (US). XIANG, Yi, Bin; 821 Main Street, Acton, MA 01720 (US).  <b>(74) Agents:</b> LUDWIG, S., Peter et al.; Darby & Darby P.C., 805 Third Avenue, New York, NY 10022-7513 (US).		<b>(81) Designated States:</b> CA, GD, HR, ID, IN, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> TRIAZINE ANTIVIRAL COMPOUNDS  <b>(57) Abstract</b>  The present invention provides pharmaceutical formulations comprising 1,3,5-triazine derivatives. The compounds and formulations of the present invention exhibit a range of activities, including antiviral and antibiotic activities, and the formulations may be used, alone or in combination, as a method of treating a patient in need of antiviral and/or antibiotic therapy. The triazine derivatives of the present invention bind to and inhibit functional nucleic acids, and hence, have broad applicability in the treatment of conditions associated with DNA and RNA viruses.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

5

## TRIAZINE ANTIVIRAL COMPOUNDS

### Field of the Invention

10                    This invention pertains to the interaction between nucleic acids and 1,3,5-triazine derivatives, their use as inhibitors of the replication of the Hepatitis B virus (HBV), and their use in the treatment of viral hepatitis caused by HBV.

### Background of the Invention

15                    There is a great medical need for novel therapeutic drugs to treat viral infections. Approximately 300,000 Americans become infected annually with hepatitis B virus (HBV). Currently available drugs for the treatment of HBV have limited efficacy and have not exhibited lasting effects, such that virus titres rapidly increase following the termination of drug treatment. In addition, some classes of  
20                    drugs, such as nucleoside analogs that inhibit the viral polymerase, often become ineffective due to the rapid appearance of resistant viral strains. For these reasons, it has become apparent that combination therapies utilizing multiple drugs that function by different mechanisms are likely to be more successful in the treatment of viral infections.

25                    Small molecules can bind RNA with high affinity and specificity and can block essential functions of the bound RNA. Examples of such molecules are

- 2 -

5 antibiotics such as erythromycin and aminoglycosides. The first suggestion that  
some antibiotic translation inhibitors interact specifically with RNA came from the  
genetic mapping of resistance to kanamycin and gentamicin to the methylation of  
16S RNA (Thompson et al., *Mol. Gen. Genet.* **201**:168, 1985). Erythromycin binds  
to bacterial RNA and releases peptidyl-tRNA and mRNA (Menninger et al.,  
10 *Mol. Gen. Genet.* **243**:225, 1994). 2-DOS-containing aminoglycosides bind  
specifically to the structures of HIV RNA known as the RRE, block binding of the  
HIV Rev protein to this RNA, and thereby inhibit HIV replication in tissue culture  
cells (Zapp et al., *Cell* **74**:969, 1993). In addition, although aminoglycosides have  
long been developed as translation inhibitors, they were only recently shown to bind  
15 to rRNA in the absence of proteins (Purohit and Stern, *Nature* **370**:659, 1994).  
Hygromycin B inhibits coronaviral RNA synthesis and is thought to do so by  
binding to the viral RNA and blocking specifically the translation of viral RNA  
(Macintyre et al., *Antimicrob. Agents Chemother.* **35**:2630, 1991). Therefore,  
compounds that bind to functionally important regions of nucleic acids of viruses  
20 and microorganisms may be useful as inhibitors of replication or other functions,  
*i.e.*, as antiviral agents and antibiotics.

The present invention pertains specifically to a novel class of drugs  
comprising RNA ligands that alter the function(s) of their target RNAs. This class  
of compounds comprises substituted 1,3,5-triazine derivatives that specifically  
25 recognize an essential and multifunctional RNA structure of the HBV pregenomic  
RNA known as the encapsidation signal ( $\epsilon$ RNA). It has been unexpectedly found  
that this class of compounds can function as inhibitors of HBV replication.  $\epsilon$ RNA,  
shown in Figure 2A, consists of a short sequence that folds into a stem-loop  
structure interrupted by a 6 nucleotide bulge.  $\epsilon$ RNA, which is contained within the  
open reading frame encoding the HBV precore protein, is also required for various  
30 steps of the HBV viral replication cycle, including encapsidation of the pregenome  
into viral particles and initiation of minus strand DNA synthesis.  $\epsilon$ RNA may also  
play a role in folding and activation of the HBV-encoded polymerase.



- 3 -

5                   Derivatives of melamine, 1,3,5-triazine-2,4,6-triamine, have been reported in the literature as suitable for various uses. For example, 2,4,6-tris(dimethylamino)-1,3,5-triazine is an antitumor agent known as Altretamine®, used in the treatment of ovarian cancer (*Cancer* 71; 4 Suppl.: 1559, 1993). Similarly, Larvadex (N-cyclopropyl-1,3,5-triazine-2,4,6-triamine) has been used as  
10 an additive to animal feed stock to control house fly infestation in poultry houses (*Poult. Sci.* 62(12): 2371, 1983).

                  Further, Patel *et al.* (*J. Inst. Chemists (India)*, 57, 1985) report a number of derivatives of 2-aryl amino-4-(4-methoxy anilino)-6-(4-chlorophenyl/phenyl hydrazido)-1,3,5-triazine having anti-bacterial activity, without  
15 data in support of this conclusion and absent any suggestion that the compounds could be used as antiviral agents.

                  U.S. Patent No. 5,225,405 to Paramelle *et al.* refers to 4,6-bis-allylamino-1,3,5-triazin-2-yl derivatives which reverse acquired resistance to anti-cancer and anti-malarial agents. Paramelle *et al.* state that the disclosed triazine  
20 derivatives, when administered at the same time with a cytotoxic agent, reduce or completely suppress multidrug resistance. The triazine compounds presumably act by inhibiting the action of an inducible membrane protein that normally functions to increase the efflux of the cytotoxic agent, thereby reducing its intracellular concentration. Paramelle *et al.* are silent as to the use of the triazine derivatives as  
25 antiviral agents.

                  U.S. Patent No. 4,508,898 to Ogilvie relates to nucleoside analogs that have a 1,3,5-triazine moiety, wherein the analog compounds exhibited antiviral activity. The compounds of Ogilvie, however, do not comprise 2,4,6-triamino-1,3,5-triazine derivatives, but rather, are N-substituted purine and pyrimidine  
30 compounds.

                  European Patent Application No. 172 608 to Kim *et al.* relates to 1,3,5-triazine derivatives that exhibit anti-ulcer, anti-inflammatory and anti-

- 4 -

5 depressant activities. However, Kim *et al.* fail to suggest that the disclosed triazine derivatives can be used as antiviral agents.

Golankiewicz, *et al.* (*J. Med. Chem.* 38: 3558, 1995) report the isolation of several 1,3,5-triazine derivatives having antiviral activity. However, the derivatives were limited to imidazo-[1,5- $\alpha$ ]-1,3,5-triazine derivatives, with  
10 special emphasis on thio- and benzyl-substituted derivatives.

Kreutzberger *et al.* (*Arzneim.-Forsch./Drug. Res.* 36 (I)(4): 626, 1986) relates to aliphatically substituted chlorodihexylamino-1,3,5-triazines having antiviral activity. However, these compounds are structurally different from the compounds of the present invention.

15 WO 97/20825 discloses the isolation of various 1,3,5-triazine derivatives structurally distinct from those of the present invention. There is no suggestion that the triazine derivatives can be used as antiviral agents, but rather, that the compounds have utility as herbicides, insecticides, miticides, and bactericides.

20 U.S. Patent No. 4,254,122 to Brown relates to 6-acylaminotetrahydro-1,3,5-triazine-2,4-dione derivatives that exhibit analgesic activities. The disclosed uses of the compound include their use as antiinflammatory agents and as inhibitors of prostaglandin synthetase.


Further, European Patent Application No. 795 549 to Gluzman *et al.*  
25 refers to bis-aryloxy(amino)-triazinyl-oxy(amino)aryl derivatives as antiviral agents. However, unlike the compounds of the present invention, the compounds of Gluzman *et al.* are dimers, linked by bicyclic or heterocyclic substituted moieties, and Gluzman *et al.* fails to suggest the use of the monomers as therapeutic compounds and/or compositions.

30 Thus, there is a need in the art for the identification of compounds that bind to functional viral nucleic acids, thereby inhibiting viral replication, and for specific antiviral agents that inhibit the replication of Hepatitis B virus. Such antiviral agents would be useful in the treatment of viral hepatitis caused by HBV.


- 5 -

### Summary of the Invention

The present invention provides methods for inhibiting viral and/or microbial replication, preventing or treating viral and/or microbial infection, and pharmaceutical formulations for use in such methods comprising a compound of the formulae IA



or IB



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, non-aromatic heterocyclic, fused or polycyclic ring, and aryloxy;

wherein said alkyl, alkenyl or alkynyl is optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino,

- 6 -

5 carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl or alkenyl; and

wherein said aryl, aryloxy, heteroaryl, non-aromatic heterocyclic, or fused or polycyclic ring is optionally substituted by one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl or alkynyl;

or wherein R<sup>1</sup> and R<sup>2</sup> together, R<sup>3</sup> and R<sup>4</sup> together, or R<sup>5</sup> and R<sup>6</sup> together, optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring, said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

or wherein R<sup>7</sup> and R<sup>8</sup> together optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring wherein said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic and fused or polycyclic ring are optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl, with the proviso that when R<sup>7</sup> and R<sup>8</sup> together form a fused or polycyclic ring, the moiety of the fused or polycyclic ring that binds with N is non-aromatic;

and pharmaceutically acceptable salts thereof;

and a pharmaceutically acceptable carrier or diluent.

Further, it is an object of the present invention to provide antiviral and antibiotic formulations comprising one or more compounds represented by the

- 7 -

5 formulae set forth above and pharmaceutically acceptable salts thereof, and a  
pharmaceutically acceptable carrier or diluent, and methods of administering such  
formulations to a patient in need of antiviral and/or antibacterial therapy. It is also  
an object of the present invention to provide a method of detecting a target nucleic  
acid by contacting the target nucleic acid with at least one compound of the  
10 formulae set forth above and pharmaceutically acceptable salts thereof, and  
monitoring an interaction between the target nucleic acid and the at least one  
compound of the formulae set forth above.

#### **Brief Description of the Drawings**

15 Figures 1A-1K are graphic illustrations of preferred 1,3,5-triazine  
compounds of the present invention.

Figure 2A is a schematic illustration of the HBV pregenomic  
sequence that corresponds to the encapsidation signal ( $\epsilon$ RNA).

20 Figure 2B is a schematic illustration of the target RNAs used in the  
method of the present invention.

Figure 3 is a graphical illustration of the reduction in virus  
production when compound 5 and the antiviral drug 2'-deoxy-3'-thiacytidine were  
tested at various concentrations alone and in combinations in an antiviral assay  
using 2.2.15 cells producing HBV.

25 Figures 4A, 4B, and 4C are graphical illustrations of the percentage  
viral reduction when compounds 5 and antiviral drug 2'-deoxy-3'-thiacytidine were  
tested in combination, with molar ratios of 1:15, 1:5, and 1:1.5, respectively.

Figures 5A and 5B are photographic and schematic representations,  
respectively, of the results of the reaction of rRNA,  $\epsilon$ RNA, RNase, and compound  
30 5 at concentrations ranging from 0 and 200  $\mu$ M.

Figures 6A and 6B are graphical and schematic illustrations,  
respectively, of the change in the melting temperature ( $t_m$ ) of  $\epsilon$ RNA in the presence  
and absence of compound 6.

5

**Detailed Description of the Invention**

All patent applications, patents, and literature references cited in this specification are hereby incorporated by reference in their entirety. In case of conflict, the present description, including definitions, will prevail.

10

**Definitions**

As used herein, the term "ligand" refers to an agent that binds a target RNA. The agent may bind the target RNA when the target RNA is in a native or alternative conformation, or when it is partially or totally unfolded or denatured. According to the present invention, a ligand can be an agent that binds anywhere on the target RNA. Therefore, the ligands of the present invention encompass agents that in and of themselves may have no apparent biological function beyond their ability to bind to the target RNA.

As used herein, the term "test ligand" refers to an agent, comprising a compound, molecule, or complex, which is being tested for its ability to bind to a target RNA.

As used herein, the term "target RNA" refers to a RNA sequence for which identification of a ligand or binding partner is desired. Target RNAs include without limitation sequences known or believed to be involved in the etiology of a given disease, condition or pathophysiological state, or in the regulation of physiological function. Target RNAs may be derived from any living organism, such as a vertebrate, particularly a mammal and even more particularly a human, or from a virus, bacterium, fungus, protozoan, parasite or bacteriophage. Target RNAs may comprise wild type sequences, or, alternatively, mutant or variant sequences, including those with altered stability, activity, or other variant properties, or hybrid sequences to which heterologous sequences have been added. Furthermore, target RNA as used herein includes RNA that has been chemically

- 9 -

5 modified, such as, for example, by conjugation of biotin, peptides, modified bases, fluorescent molecules, and the like.

Target RNA sequences for use in the present invention are typically between about 5 and about 500 nt, preferably between about 30 and about 100 nt, and most preferably about 50 nt. Target RNAs may be isolated from native  
10 sources, or, more preferably, are synthesized *in vitro* using conventional polymerase-directed cell-free systems such as those employing T7 RNA polymerase. In a preferred embodiment, the target RNA is HBV  $\epsilon$ RNA. Figure 2A shows the actual portion of the HBV pregenomic sequence that corresponds to the encapsidation signal ( $\epsilon$ RNA). Figure 2B shows the  $\epsilon$ RNA sequence used as  
15 target RNA in the method of the present invention, and the RRE RNA target used for specificity tests. The target RNA was prepared by *in vitro* transcription of a linearized plasmid containing the cloned target sequence, with bacteriophage SP6 RNA polymerase in the presence of  $\alpha$ P<sup>32</sup>-UTP to obtain radiolabeled target RNA using methods known to those of ordinary skill in the art.

20 As used herein, the term "treatment" with regard to a viral or microbial infection includes preventing, retarding, and/or reducing a disease, pathological condition or one or more symptoms thereof, in vertebrates, *e.g.*, birds, and mammals, particularly humans. In the case of HBV, the altering or inhibiting any of the following processes can be considered very useful for the treatment of  
25 infection mediated by HBV. They are:

1. Symptoms associated with acute hepatitis, including the onset of the prodromal phase, which is accompanied by anorexia, malaise, nausea and vomiting, and often fever, and which is followed in the icteric phase by the occurrence of urticarial eruptions, arthralgias and jaundice;
- 30 2. Elevations in serum aminotransferase and urinary bile levels prior to and during the onset of maximal jaundice;
3. Low-normal white blood cell counts, and appearance on blood smears of atypical lymphocytes; or

- 10 -

5                   4. Symptoms associated with chronic hepatitis infection, including viremia, seroconversion, liver cancer, and cirrhosis.

                  According to the present invention, treatment constitutes any improvement in one or more clinical or histological symptoms or diagnostic markers observed by the attending physician or determined by quantitative or  
10                   semiquantitative techniques. Non-limiting examples of appropriate techniques include analysis of blood and urine.

                  As used herein, "inhibition of replication" refers to any detectable reduction in replication or growth of virus, bacteria or fungi, e.g. between about 1 % and about 100 % reduction, preferably between about 5 % and about 100 %  
15                   reduction, and more preferably between about 10 % and about 100 % reduction. The skilled artisan will appreciate that any reduction in viral replication, bacterial or fungal growth is significant where it is approximately equal to or greater than that which is observed for known inhibitors of viral replication, bacterial or fungal growth.

20                   As used herein, the terms "antibiotic", "antibacterial" and "antifungal" refer to any compound that inhibits growth of or destroys microorganisms, bacteria, or fungi.

                  As used herein, the term "aryl" means an aromatic carbocyclic ring system having a single radical containing about 6 to about 10 carbon atoms. An  
25                   aryl group may be a fused or polycyclic ring system. Exemplary aryl groups include phenyl or naphthyl.

                  As used herein, the term "aryloxy" means an O-aryl group. An aryloxy group is optionally substituted on the aryl moiety of the aryloxy. Suitable substituents include halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy,  
30                   alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl.



- 11 -

5                   As used herein, the term "alkyl" means a straight or branched saturated hydrocarbon group. Preferred alkyl groups include those having from 1-12 carbon atoms.

                  As used herein, the term "cycloalkyl" means a non-aromatic monocyclic or fused or polycyclic ring system of about 3 to about 10 ring carbon atoms. Optionally one or more of the ring carbon atoms of the cycloalkyl may be replaced by a heteroatom, such as nitrogen, oxygen or sulfur. Exemplary cycloalkyl groups include cyclohexyl.


10

                  As used herein, the term "cycloalkenyl" means a non-aromatic monocyclic or fused or polycyclic ring system containing a carbon-carbon double bond and having about 3 to about 10 ring carbon atoms. Optionally one or more of the ring carbon atoms of the cycloalkyl may be replaced by a heteroatom, such as nitrogen, oxygen or sulfur.

15

                  As used herein, the term "carbonyl" or "carbonyl moiety" refers to any chemical moiety comprising a carbonyl functional group, *e.g.*, a ketone, aldehyde, carboxylic acid, acid halide, amide, peptide, anhydride and ester. As used herein, when a ring structure is described as substituted with a carbonyl group, the ring carbon atom is replaced by the group

20



                  As used herein, the term "heteroatom" includes nitrogen, oxygen and sulfur, as well as any atom other than a carbon.


25

                  As used herein, the term "ring system" refers to an aromatic or non-aromatic carbocyclic compound, in which one or more of the ring carbon atoms may be replaced by

- 12 -


5 a heteroatom, such as nitrogen, oxygen or sulfur. The ring system may be optionally substituted by one or more halogens, C<sub>1</sub> to C<sub>12</sub> alkyl, aryl, vinyl, alkyl(aryl), vinyl(aryl) and nitro groups.

As used herein, the term "fused ring system" refers to ring systems wherein at least two adjacent carbon centers join one or more cyclic structures. A  
10 fused ring system as used herein may be aromatic or non-aromatic, or may be composed of separate aromatic and non-aromatic moieties. Exemplary carbocyclic fused ring systems are represented by the formulas:



15

Exemplary fused ring systems in which one or more of the ring carbon atoms is replaced by a heteroatom include the following:




20

- 13 -

5


As used herein, the term "polycyclic ring system" refers to ring systems having two or more cyclic compounds bonded in tandem. A polycyclic ring system as used herein may be aromatic or non-aromatic, or may be composed of separate aromatic and non-aromatic moieties. An exemplary carbocyclic

10 polycyclic ring system is represented by the formula



An exemplary polycyclic ring system in which one or more of the ring carbon atoms is replaced by a heteroatom include the following:

15



Additionally, fused or polycyclic ring systems may optionally be substituted by one or more halogens, C<sub>1</sub> to C<sub>12</sub> alkyl, aryl, vinyl, alkyl(aryl), vinyl(aryl) and nitro groups.

20

As used herein, the term "heteroaryl" means an about 5 to 10-membered aromatic monocyclic or fused or polycyclic ring system having a single radical in which one or more of the carbon atoms in the ring system is other than carbon, for example, nitrogen, oxygen or sulfur. An exemplary heteroaryl group is pyridine. An exemplary fused or polycyclic heteroaryl group is indole.

25

- 14 -

5                   As used herein, the term "heterocyclyl" or "heterocyclic" means an aromatic or non-aromatic about 5 to about 10-membered monocyclic or fused or polycyclic ring system in which one or more of the carbon atoms in the ring system is other than carbon, for example, nitrogen, oxygen or sulfur. A heterocyclyl group may be a fused or polycyclic ring system. Exemplary heterocyclyl groups  
10 include piperidine, morpholino, and azepanyl.

                  As used herein, the term "primary, secondary, or tertiary amine" refers to amine compounds having one, two, or three functional groups, respectively. Suitable functional groups include halogens, amines, C<sub>1</sub> to C<sub>12</sub> alkyl groups, aryl, vinyl, alkyl(aryl), vinyl(aryl) and nitro groups.

15                   As used herein, the term "amide" refers to groups having the amide functional group -C(O)NH-, wherein alkyl, alkenyl or alkynyl groups may be bonded to the C or N atom of the amide group.


                  As used herein, the term "high affinity" refers to a compound that binds tightly to its target. Preferred compounds of the present invention will exhibit  
20 an IC<sub>50</sub> at or below 50 μM, preferably at or below 10 μM, and more preferably at or below 1 μM.

                  As used herein, the term "specificity" refers to a compound having either high or low affinity for its target. A highly specific compound will be unaffected by competitor RNA and will not have an effect on a heterologous assay  
25 using a different target, independent of the compound's affinity. In a preferred embodiment, compounds of the present invention will exhibit no activity or at least a 5-fold higher IC<sub>50</sub> value in a heterologous assay than in a specific assay.

                  The present invention provides methods for inhibiting the replication of viruses and microorganisms and for preventing or treating viral or microbial  
30 infection, which comprise administering 1,3,5-triazine compounds and pharmaceutically acceptable salts thereof. The compounds bind Hepatitis B virus (HBV) eRNA with high affinity and specificity, and thereby alter its function. The


- 15 -

5 formulations of the invention comprise triazine derivatives represented by the formulae IA



or IB

10



15

wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, non-  
20 aromatic heterocyclic, fused or polycyclic ring and aryloxy;

wherein said alkyl, alkenyl or alkynyl is optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano,  
25 cycloalkyl, alkenyl, cycloalkenyl or alkynyl; and

- 16 -

5                                    wherein said aryl, aryloxy, heteroaryl, non-aromatic heterocyclic or fused or polycyclic ring is optionally substituted by one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl or  
10                                    alkynyl;

                                  or wherein R<sup>1</sup> and R<sup>2</sup> together, R<sup>3</sup> and R<sup>4</sup> together, or R<sup>5</sup> and R<sup>6</sup> together, optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring, said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring optionally substituted with one  
15                                    or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

                                  or wherein R<sup>7</sup> and R<sup>8</sup> together optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic or fused or polycyclic ring, wherein said cycloalkyl, cycloalkenyl and non-aromatic heterocyclic or fused or polycyclic ring are optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines,  
20                                    cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl, with the proviso that when R<sup>7</sup> and R<sup>8</sup> together form a fused or polycyclic ring, the moiety of the fused or polycyclic ring that binds with N is non-aromatic;



                                  and pharmaceutically acceptable salts thereof;

                                  and a pharmaceutically acceptable carrier or diluent.

30                                    In one embodiment, the invention is directed to compounds of formula IB wherein one or R<sup>1</sup> and R<sup>2</sup> is an optionally substituted aryl. In another embodiment, the invention is directed to compounds of formula IB wherein one of


- 17 -

- 5       $R^7$  or  $R^8$  is an optionally substituted aryl      Non-limiting examples of the  
compounds of the invention include:




- 18 -

5



IV




V

10




- 19 -

5



VI




VII

10

The following is a chemical process for the efficient production of triazines of the formula IA:

15




20

or IB

- 20 -

5

10



Commercially

15

available cyanuric chloride (IX) is first reacted with two equivalents of a reagent chosen from a group consisting of a primary amine and a secondary amine, to afford a singly substituted triazine of the formula (X).




Fig. IX

Fig. X

20

- 21 -

- 5 The 2-amino 4,6 di chloro-1,3,5 triazine of the formula (X) is then reacted with two equivalents of a reagent chosen from a group consisting of a primary amine and secondary amine to afford a disubstituted triazine of the formula (XI)




Fig. X

Fig. XI

- 10 The 2,4-diamino-6-chloro-1,3,5 triazine of the formula (XI) can then be reacted with two equivalents of a reagent chosen from a group consisting of a primary amine, secondary amine, and hydrazine. In the case of a primary and secondary amine, the reaction affords a trisubstituted triazine of the formula (XII).




Fig. XI

Fig. XII

In the case of hydrazine, condensation reaction forms compound (XIII).

- 22 -




Fig. XI

Fig. XIII

5

Compound (XII) can then be further reacted with a reagent chosen from a group consisting of aldehydes and ketones to afford 2,4,6- substituted-1,3,5-triazines of the formula (XIV).




Fig. XIII

Fig. XIV

10

### Detailed Description of the Preferred Embodiments

The following is a chemical process for the efficient production of triazine derivatives:

Example 1: 2-fluorobenzaldehyde N-[4-(benzylamino)-6-(tert-butylamino)-1,3,5-triazin-2-yl]hydrazone (compound 93).

15


- 23 -

5

10

15


20



- 24 -

## 5 Step 1:

To a stirred solution of cyanuric chloride in DME at -30 °C was added a solution of 2 equivalents of benzylamine in water in dropwise fashion. The mixture was then stirred for 3 hrs. at the reduced temperature, after which time the mixture was warmed to room temperature and washed sequentially with saturated sodium bicarbonate and water, dried over magnesium sulfate, and reduced in vacuo. The product was used in the next step without purification.




- 25 -

## 5      Step 2:

To a stirred solution of the product of step 1 in dichloromethane at room temperature was added 2 equivalents of tert-butyl amine in dropwise fashion. The solution was stirred for 12 hrs., after which time the mixture was then washed sequentially with 0.1 M hydrochloric acid, water, and saturated brine, dried over sodium sulfate, reduced in vacuo and recrystallized from hexane:ethyl acetate (4:1) to afford the starting material of step 2.

10


15



- 26 -

## 5 Step 3:

To a solution of the product of step 2 in dichloromethane was added 2 equivalents of hydrazine in water in dropwise fashion. The solution was then heated to 60°C for a period of 12 hrs., after which time the mixture was cooled to room temperature, washed sequentially with brine and water, dried over sodium sulfate and reduced in vacuo to a residue which was then recrystallized from 10 hexane:ethyl acetate (4:1) to afford the starting material of step 4.




## 15 Step 4:

To a stirred solution of the product of step 3 in toluene in a round bottom flask was added one equivalent of 2-fluorobenzaldehyde. The flask was then fitted with a Dean Stark trap and refluxed to azeotropically remove water.



- 27 -

- 5 After which time the removal of water was complete, the solution was cooled to room temperature and the solvent was removed in vacuo. The residue was then recrystallized from hexane:ethyl acetate (5:1) to afford compound 93.




- 28 -

5     Example 2: 2-hydroxybenzaldehyde N-[4-(benzylamino)-6-(diethylamino)-1,3,5-triazin-2-yl]hydrazone (compound 155).

Step 1:

10            To a stirred solution of cyanuric chloride in DME at -30 °C was added a solution of 2 equivalents of benzylamine in water in dropwise fashion. The mixture was then stirred for 3 hrs. at the reduced temperature, after which time the mixture was warmed to room temperature and washed sequentially with saturated sodium bicarbonate and water, dried over magnesium sulfate and reduced in vacuo. The product was used in the next step without purification.

15




Step 2:

25            To a stirred solution of the product of step 1 in dichloromethane at room temperature was added 2 equivalents of diethyl amine in dropwise fashion. The solution was stirred for 12 hrs., after which time the mixture was then washed

- 29 -

- 5 sequentially with 0.1 M hydrochloric acid, water and saturated brine, dried over sodium sulfate, reduced in vacuo and recrystallized from hexane:ethyl acetate (4:1) to afford the starting material of step 3.


10



- 30 -

## 5 Step 3:

To a solution of the product of step 2 in dichloromethane was added 2 equivalents of hydrazine in water in dropwise fashion. The solution was then heated to 60°C for a period of 12 hrs., after which time the mixture was cooled to room temperature, washed sequentially with brine and water, dried over sodium sulfate and reduced in vacuo to a residue which was then recrystallized from 10 hexane:ethyl acetate (4:1) to afford the starting material of Step 4.




## 15 Step 4:

To a stirred solution of the product of step 3 in toluene in a round bottom flask was added one equivalent of salicylaldehyde. The flask was then fitted with a Dean Stark trap and refluxed to azeotropically remove water. After

- 31 -

5 which time the removal of water was complete, the solution was cooled to room temperature and the solvent was removed in vacuo.




10

The residue was then recrystallized from hexane:ethyl acetate (5:1) to afford compound 155:

15

- 32 -

5



#### Determination of Antiviral or Antimicrobial Activity

##### In Vitro Assay

One assay used in determining the activity of the compounds of the present invention is disclosed in copending U.S. Patent Application Serial No. 08/709,342, filed September 6, 1996, the disclosure of which is hereby incorporated herein by reference in its entirety. The assay detects the interaction between target RNA molecule and a test ligand by measuring the ligand's ability to inhibit hybridization between the RNA and a specific complementary oligonucleotide. In one embodiment, the target RNA is radiolabelled and the oligonucleotide is labeled with biotin; in this case, the extent of hybridization is determined by measuring the amount of radiolabeled RNA detected with a streptavidin/biotin-based capture system.

- 33 -

5                   Ligands that exhibit inhibitory activity in a primary *in vitro* assay are then titrated and tested in secondary assays to eliminate false positives.

Concentrations of test compounds of between about 0.1  $\mu$ M and about 200  $\mu$ M are assayed for inhibition using radiolabeled  $\epsilon$ RNA in the absence or presence of a large excess of an unlabeled non-specific RNA competitor (such as, *e.g.*, ribosomal  
10   RNA, rRNA). Compounds that bind  $\epsilon$ RNA non-specifically are competitively displaced by the rRNA, resulting in reduced or no inhibition of hybridization between the radiolabeled  $\epsilon$ RNA and the biotinylated oligonucleotide. Conversely, the inhibitory activity of compounds that specifically bind  $\epsilon$ RNA is not affected by the presence of the competitor rRNA.

15                   Specificity is further tested in a third assay in which the radiolabeled RNA is the HIV derived RRE RNA and the biotinylated oligonucleotide is complementary to RRE. Compounds with high specific activity for  $\epsilon$ RNA are not expected to be inhibitory in the RRE RNA based assay. The specificity of the ligand for  $\epsilon$ RNA is expressed as the ratio between the  $IC_{50}$  value obtained with the  
20   specific target RNA and the  $IC_{50}$  value obtained with a heterologous target (RRE RNA), both in the presence of competitor RNA.

The substituted 2,4,6-triamino-1,3,5-triazine derivatives of the present invention preferably exhibit  $IC_{50}$  values for their interaction with  $\epsilon$ RNA at or below 300  $\mu$ M, more preferably at or below 50  $\mu$ M, and most preferably at or  
25   below 5  $\mu$ M, in the presence of an excess of non-specific competitor rRNA.

- 34 -

5

Bioassay

Ligands that exhibit high affinity and specificity for  $\epsilon$ RNA as determined above are tested for their ability to inhibit HBV replication. Several cell lines and cell culture assays have been developed to identify potential therapeutics effective against chronic HBV infection. One of these cell lines, 2.2.15, has been used in a standardized assay that has repeatedly proven to be an accurate model of chronic cellular HBV replication and a predictive model of antiviral response for chronic hepadnaviral infection *in vivo*. 2.2.15 cells contain copies of the complete HBV genome integrated into the cell's genome, and 2.2.15 cells are not susceptible to infection by HBV. (Sells, M. A. *et al.*, J. Virol. 62 (8): 2836-2844 (1988)).

10 These cells express HBV genes producing complete HBV particles capable of infecting chimpanzees (Acs, G. *et al.*, Proc. Natl. Acad. Sci. USA 84 (13): 4641-4644 (1987)). Briefly, 2.2.15 cells are cultured and pretreated with a test compound for about 9 days, at which point the media are harvested and subjected to dot-blot hybridization to detect HBV virion DNA.

20

Further, the antiviral effect of any compound must be measured against its toxicity. Cytotoxicity is measured by the neutral red uptake method using the same cells used for antiviral activity.

25

It will be understood that the skilled artisan can employ any assay suitable for assessing the inhibitory potency of a compound to practice the present invention without undue experimentation. See, *e.g.*, J. Sambrook and T. Maniatis, "Molecular Cloning: A Laboratory Manual", Cold Spring Harbor Laboratories



- 35 -

5 Press, Cold Spring Harbor, NY (1989), "Current Protocols in Molecular Biology",  
Ed. F.M Ausubel, *et al.*, J. Wiley & Sons, Inc. 1997, D. Leland, "Clinical  
Virology", W.B. Saunders Co., 1996, and "Cells: A Laboratory Manual", Ed.  
D.L. Spector, *et al.* Cold Spring Harbor Laboratory Press, Cold Spring Harbor,  
New York, 1997.

10 The compounds of the present invention are believed to inhibit viral  
or microbial replication by binding to functionally important viral or microbial  
nucleic acids. The nucleic acids may be RNA or DNA, may form part of the  
genome of the virus or microorganism or an intermediate thereof, or may represent  
expressed mRNA species or any other functional nucleic acid unique to the  
15 replication or function of the virus or microorganism.

Compounds of the present invention inhibit hybridization of the  
oligonucleotide probe in the assay described in copending application U.S. Patent  
Application Serial No. 08/709,342, filed September 6, 1996. This indicates that the  
compounds interact with the RNA target by stabilizing the RNA structure, thereby  
20 inhibiting the formation of hybrids between the RNA target and the complementary  
oligonucleotide present in the assay.

Supplementary evidence in support of the contention that triazine  
compounds interact with the RNA target came from the observation that compound  
5 protects specific positions of RNA from digestion with RNases. RNase protection  
was assessed by incubating an end-labeled RNA with the amount of RNase needed  
25 to yield a ladder representing single cut events at each possible cleavage site in the

- 36 -

5 RNA structure (based upon a gel electrophoresis analysis). The presence of a ligand bound to the RNA renders the binding site inaccessible to the RNases and therefore, the binding site will be shown by the disappearance or weakening of specific bands in the digestion pattern (Figure 5).

Further, the RNA binding ability of the triazine compounds was  
10 demonstrated by preparing an RNA target containing only the bulge region of  $\epsilon$ RNA. The melting temperature of the RNA was determined by monitoring the change in optical density at 260 nm with increasing temperature. Upon addition of 10  $\mu$ M of triazine compound, the melting temperature was shifted, which indicates that the RNA structure is stabilized by the ligand (Figure 6).

15 Thus, these experiments demonstrate that the triazine compounds interact with the bulge region of  $\epsilon$ RNA by nucleic acid binding.

The methods and formulations of the invention can be used to inhibit replication and/or prevent or treat infection caused by: hepatitis B virus, hepatitis C virus, herpes simplex virus, types 1 and 2, varicella-zoster, cytomegalovirus,  
20 Epstein-Barr virus, polyomavirus, papillomavirus, parvovirus, vaccinia virus, molluscum contagiosum, Marburg and Ebola viruses, influenza A and B, measles, mumps, respiratory syncytial virus, poliovirus, coxsackie virus A and B, rhinovirus, rotavirus, human immunodeficiency virus, types 1 and 2, rabies, rubella, and equine encephalitis. DNA and RNA viruses encompassed by the  
25 invention, include, without limitation, viruses belonging to the viral families adenoviridae, hepadnaviridae, herpesviridae, papovaviridae, parvoviridae,

- 37 -

5       poxviridae, arenaviridae, bunyaviridae, flaviviridae, orthomyxoviridae,  
paramyxoviridae, picornaviridae, reoviridae, retroviridae and rhabdoviridae.

          Accordingly, compounds and formulations of the present invention  
can be used in the prevention and treatment of viral hepatitis caused by HBV, as  
well as in the prevention and treatment of disease conditions associated with other  
10       DNA and RNA viruses. Such conditions include, but are not limited to: upper and  
lower respiratory tract infections, ocular infections, gastroenteritis, cystitis, and  
complications arising in transplant recipients, each of which is associated with  
adenoviruses; treatment of immunocompromised individuals, wherein a  
compromised immune system is associated with cytomegalovirus; infectious  
15       mononucleosis, associated with Epstein-Barr virus; chickenpox and shingles, which  
are associated with varicella-zoster; oral, genital and skin lesions, as well as various  
dermatological anomalies associated with herpesvirus, papillomavirus and  
polyomavirus; smallpox, as well as complications arising from smallpox  
vaccinations, including allergic rash, progressive vaccinia and postvaccinial  
20       encephalitis, each of which being associated with variola, molluscum and  
contagiosum; hemorrhagic fever and aseptic meningitis, which are associated with  
lassa fever virus, lymphocytic choriomeningitis virus and other arenaviruses; upper  
and lower respiratory infections, serious acute respiratory tract illness and  
pneumonia associated with influenza A, B, and C; measles, mumps, and  
25       parainfluenza, associated with paramyxoviruses; enteric, neuromuscular and central  
nervous system infections associated with picornaviruses; acquired immune

- 38 -

5        deficiency syndrome (AIDS) and associated infections and clinical syndromes  
associated with HIV infection; and encephalitis and measles associated with  
togaviruses.

10            Moreover, given the functional similarities of the compounds of the  
present invention and other RNA binding molecules, *e.g.*, neomycin, erythromycin  
and aminoglycosides, the compounds and compositions of the present invention can  
be used as antibiotic therapies. Antibiotics are used in the treatment of infectious  
diseases in plants, animals and man, and may kill and/or inhibit the growth of  
microorganisms. Therefore, the compounds and compositions of the present  
invention may be used as bacteriocidal, bacteriostatic and broad-spectrum antibiotic  
15        agents.

#### **Pharmaceutical Formulations**

20            The compositions of the present invention can be administered in  
dosages and by techniques well known to those skilled in the medical, veterinary,  
and agricultural arts taking into consideration such factors as the age, sex, weight,  
species and condition of the particular patient, and the route of administration. The  
compositions of the present invention can be administered alone, or can be co-  
administered or sequentially administered with additional, non-triazine based  
antiviral agents, such as, *e.g.*, Acyclovir,  $\alpha$ -interferon, ribavirin, and various  
25        protease inhibitors, *e.g.*, ritonavir, indinavir, saquinavir, and/or additional non-

- 39 -

5 triazine based antibiotics, such as, *e.g.*, erythromycin, gentamycin, nanamycin, and streptomycin.

The effect of simultaneous administration of triazine compound 5 and 2'-deoxy-3'-thiacytidine, a well known nucleoside analog having antiviral activity, was examined in order to assess the interaction between the two therapies. The results show a strong synergistic interaction when the two drugs are administered together at a 1:15 molar ratio (3TC: triazine; Figure 4A). The antiviral activity observed with the combination drug treatment is larger than the additive activity expected based on the activities of each compound administered alone. A synergistic response allows the use of lower amounts of each drug in combination therapies, which reduces the toxicity and secondary risks. Moreover, combination therapies with drugs acting on different targets should reduce the probability that drug resistant viral strains will develop.

Moreover, the formulations of the present invention can be administered in a formulation suitable for the manner of administration, including but not limited to liquid preparations for mucosal administration, *e.g.*, oral, nasal, anal, vaginal, peroral, intragastric administration and the like, such as solutions, suspensions, syrups, elixirs. Further, liquid preparations for administration of the compositions of the present invention for parenteral, subcutaneous, intradermal, intramuscular, intravenous administrations, and the like, such as sterile solutions, suspensions or emulsions, *e.g.*, for administration by injection, can be formulated without undue experimentation. Oral administration is presently preferred.

- 40 -

5                   In order for a composition to be administered to an animal or human, and for any particular method of administration, it is preferred to determine the toxicity, such as by determining the lethal dose (LD) and LD<sub>50</sub> in a suitable animal model, *e.g.*, mouse; the dosage of the composition(s), and the concentration of components in the composition; and the timing of administration in order to  
10                   maximize the antiviral and/or antimicrobial response. Such factors can be determined without undue experimentation by such methods as titrations and analysis of sera for antibodies or antigens, *e.g.*, by ELISA and/or EFFIT analysis. Such determinations do not require undue experimentation from the knowledge of the skilled artisan, the present disclosure and the documents cited herein.

15                   The formulations can be administered in a pharmaceutically effective amount, an antiviral effective amount and/or in an antimicrobial effective amount, taking into account such factors as the relative activity and toxicity for the target indication, *e.g.*, antiviral activity and/or antimicrobial activity, as well as the route of administration, and the age, sex, weight, species and condition of the particular  
20                   patient.

                  The compositions of the present invention can be solutions, suspensions, emulsions, syrups, elixirs, capsules, tablets, and the like. The compositions may contain a suitable carrier, diluent, or excipient, such as sterile water, physiological saline, glucose, or the like. Moreover, the compositions can  
25                   also be lyophilized, and/or may contain auxiliary substances, such as wetting or emulsifying agents, pH buffering agents, adjuvants, gelling or viscosity enhancing

- 41 -

5 additives, preservatives, flavoring agents, colors, and the like, depending upon the route of administration and the preparation desired. Standard texts, such as "Remington's Pharmaceutical Science", 17th Ed., 1985, incorporated herein by reference, may be consulted to prepare suitable preparations, without undue experimentation.

10 Suitable dosages for compositions of the present invention can be determined without undue experimentation based upon the Examples provided below, and the documents cited herein. For instance, suitable dosages of antiviral and/or antibiotic agent in a composition can be 0.1 to 250 mg/kg/day, preferably below 100 mg/kg/day, and most preferably below 50 mg/kg/day.

15 In a further embodiment, the compounds of the present invention can be used as lead compounds to improve the antiviral and/or antibiotic activities of the compounds. This can be done by modifying certain functional groups of the compounds of the present invention based upon a recognition of the structure/activity relationship between a particular functional group in a compound  
20 and its biological activity. Such modifications include synthetic manipulation of the size, hydrophilicity, hydrophobicity, acidity and basicity of a functional group, which may inhibit or enhance the activity of a compound.

Moreover, based upon the observed binding interaction between the compounds of the present invention and nucleic acids, the compounds of the present  
25 invention can be used in a method of detecting and/or purification of nucleic acids, e.g. binding assays and affinity chromatography. For example, the compounds of

- 42 -

5 the invention can be covalently attached to a chromatographic support and a nucleic acid-containing solution can be eluted over the solid support, which will bind the target nucleic acid as the other components of the solution elute from the column in the flow-through. With regard to binding assays and affinity chromatography, reference is made to A. Fersht, Enzyme Structure and Mechanism, 2d Ed., W.H. Freeman & Co., New York, 1985, and R.K. Scopes, Protein Purification,  
10 Principles and Practice, 2d Ed., Springer-Verlag, New York, 1987, the disclosures of which are hereby incorporated herein by reference.

Additionally, the compounds of the present invention could be used as inhibitors of specific steps of the viral replication cycle in order to study the  
15 process *in vivo* or *in vitro*. In a diagnostic application, the compound could be used for detection of specific RNA species in samples by using a triazine derivative labeled with a fluorescent, immunochemical, or radioactive moiety. Further, such labeled triazine compounds could be used to study intracellular localization of the RNA target by electron-microscopy or light microscopy using tissue culture  
20 preparations.

The following examples are intended to further illustrate the invention without imposing any undue limitations thereon.

25 **Example 1: HIGH-THROUGHPUT IDENTIFICATION OF  $\epsilon$ RNA LIGANDS**

Compounds of the invention were obtained from commercial sources. Compound 32 is available from MicroSource Discovery Systems, Inc.,



- 43 -

5        Gaylordsville, Connecticut; compounds 64, 136, and 148 are available from  
ComGenex, Inc., Budapest, Hungary; compounds 103 and 123 are available from  
ChemBridge Corp., San Diego, California; and the remaining compounds are  
available from Specs and Biospecs B.V., Rijswijk, The Netherlands.

10        The high throughput assays were performed in 96-well plates  
containing eighty drugs (5  $\mu$ l, 60  $\mu$ g/ml in DMSO) distributed one per well in  
columns 2-6 and 8-12 of the 8 x 12 96-well array. The remaining 16 wells in  
columns 1 and 7 were filled with 5  $\mu$ l DMSO to serve as the control reactions. 45  
 $\mu$ l of reaction mixture containing buffer, salts and radiolabeled target RNA were  
delivered on each well of the compound-containing plates to yield a 50  $\mu$ l mixture  
15        containing 50 mM Tris-HCl, pH 7.5, 200 mM KCl, 5 mM MgCl<sub>2</sub>, 5 mM DTT, 6  
 $\mu$ g/ml test compound, and 25 nM [<sup>32</sup>P]-labeled  $\epsilon$ RNA. The reaction was started by  
addition of the biotinylated oligonucleotide, 262.104 A (5  $\mu$ l) to a final  
concentration of 100 nM. The reaction was incubated at 25 °C for 60 minutes, and  
then 5  $\mu$ l of 0.3 mg/ml SAAP (streptavidin alkaline phosphatase conjugate, Pierce,  
20        Rockford, Ill.) was added. The reaction was incubated for an additional 30 minutes  
at 25°C and filtered through 96-well format HATF nitrocellulose filters (Millipore,  
Bedford, MA) using a multiscreen vacuum manifold (Millipore, Bedford, MA).  
Subsequently, 300  $\mu$ l of wash buffer (50 mM Tris-HCl, pH 7.5, 200 mM KCl) was  
filtered through to wash the filters. The filters were dried and supplemented with  
25        20  $\mu$ l scintillation fluid (Super Mix, Wallac Oy, Turku, Finland). The amount of

- 44 -

5 hybrids formed was determined by scintillation counting of the radiolabeled  $\epsilon$ RNA retained on the filters.

The inhibitory effect of a test compound (expressed as percentage of inhibition, % Inh.) was calculated using the following formula:

$$\% \text{ Inh.} = (1 - (R_x/R_o)) \times 100$$

10 where  $R_o$  is the average retention of hybrids in the 16 identical control reactions in columns 1 and 7 and  $R_x$  is the hybrids retained in the presence of a test drug,  $x$ .

Biotinylated oligonucleotides 262.104 A (5'-biotin-TTA GGC ACA GCT TGG AGG CTT GAA CAG TG-3') and 208.92 A (5'-biotin-CGT CAT TGA CGC TGC GCC CA-3') were synthetically prepared (Oligos Etc., Willsonville,  
15 OR). These oligonucleotides are complementary to regions of the  $\epsilon$ RNA and RRE RNA targets, respectively.

A plasmid containing the target sequence following the bacteriophage SP6 promoter was used as a template for the synthesis of radiolabeled  $\epsilon$ RNA using [ $\alpha^{32}$  P]-UTP and SP6 RNA polymerase as described by the suppliers, *e.g.*, Ambion,  
20 Austin, TX, and by methods well known to those of ordinary skill in the art.

Compounds that inhibited the primary screening assay described above were individually titrated in the same assay at concentrations between 0.1  $\mu$ M and 200  $\mu$ M in the absence and presence of 60-fold excess rRNA as a non-specific competitor. A third titration was performed in a heterologous assay using  
25 radiolabeled RRE RNA as the target and the complementary biotinylated

- 45 -

5 oligonucleotide 208.92 A. From these three assays, the  $IC_{50}$  values and specificity  
for each compound was calculated.

### Results:

$IC_{50}$  values using rRNA competitor (from Example 1) and  
10 radiolabeled RRE RNA are shown in Table 1 for the following ligands. A  
compound is considered inactive if the titration course does not result in sufficient  
inhibition to estimate an  $IC_{50}$  value. Specificity is calculated as the ratio of RRE +  
rRNA  $IC_{50}$ /εRNA + rRNA  $IC_{50}$ . When an εRNA ligand was inactive in the RRE  
assay the specificity of the ligand is defined as maximal (max) since a value cannot  
15 be calculated.

Table 1.  $IC_{50}$  Values for 1,3,5-Triazine Compounds

<u>Compound</u>	<u>Specificity</u>	<u>HBV+ rRNA</u> <u><math>IC_{50}</math> (<math>\mu</math>M)</u>	<u>RRE + rRNA</u> <u><math>IC_{50}</math> (<math>\mu</math>M)</u>
1	max.	32.4	inactive
2	max.	3.2	inactive
3	max.	66.9	inactive
4	max.	16.3	inactive
5	max.	49.3	inactive
6	max.	10.4	inactive
7	max.	2.4	Inactive
8	max.	4	inactive

- 46 -

	<u>Compound</u>	<u>Specificity</u>	HBV+ rRNA <u>IC<sub>50</sub> (μM)</u>	RRE + rRNA <u>IC<sub>50</sub> (μM)</u>
5	9	132.6	6.2	inactive
	10	max.	12	inactive
	11	max.	15	inactive
	12	max.	2.6	inactive
	13	max.	4.9	inactive
10	14	max.	7.1	inactive
	15	max.	11	inactive
	16	max.	12.1	inactive
	17	max.	15.1	inactive
	18	max.	5.1	inactive
15	19	max.	9.9	inactive
	20	max.	11.7	inactive
	21	max.	14.4	inactive
	22	max.	3.6	inactive
	23	max.	5.7	inactive
20	24	max.	10	inactive
	25	max.	11.9	inactive
	26	max.	15	inactive
	27	max.	15.5	inactive
	28	max.	18.9	inactive
25	29	max.	24.8	inactive
	30	max.	25	inactive
	31	max.	30	inactive
	32	max.	32.6	inactive

- 47 -

	<u>Compound</u>	<u>Specificity</u>	HBV+ rRNA <u>IC<sub>50</sub> (μM)</u>	RRE + rRNA <u>IC<sub>50</sub> (μM)</u>
5	33	max.	37.2	inactive
	34	max.	21	inactive
	35	max.	24.9	inactive
	36	53.8	28.9	1553.8
	37	max.	30.1	inactive
10	38	max.	34.9	inactive
	39	max.	39.9	inactive
	40	max.	21.9	inactive
	41	max.	24.9	inactive
	42	max.	30	inactive
15	43	max.	30.1	inactive
	44	max.	35	inactive
	45	max.	39.9	inactive
	46	max.	23.1	inactive
	47	max.	25	inactive
20	48	max.	30	inactive
	49	max.	35	inactive
	50	max.	39.9	inactive
	51	max.	39.9	inactive
	52	max.	66	inactive
25	53	max.	70.1	inactive
	54	max.	89.9	inactive
	55	max.	100.1	inactive
	56	max.	42.1	inactive

- 48 -

	<u>Compound</u>	<u>Specificity</u>	HBV+ rRNA <u>IC<sub>50</sub> (μM)</u>	RRE + rRNA <u>IC<sub>50</sub> (μM)</u>
5	57	max.	50.1	inactive
	58	max.	200.4	inactive
	59	max.	90.1	inactive
	60	max.	100.1	inactive
	61	max.	42.6	inactive
10	62	max.	54.3	inactive
	63	max.	69	inactive
	64	max.	71.9	inactive
	65	max.	99.9	inactive
	66	max.	113.5	inactive
15	67	max.	47.7	inactive
	68	max.	33	inactive
	69	max.	70	inactive
	70	max.	80.6	inactive
	71	max.	99.9	inactive
20	72	max.	119.9	inactive
	73	max.	120	inactive
	74	max.	125.9	inactive
	75	max.	160.1	inactive
	76	max.	200	inactive
25	77	max.	120	inactive
	78	max.	133.2	inactive
	79	max.	168.9	inactive
	80	max.	200	inactive

- 49 -

	<u>Compound</u>	<u>Specificity</u>	HBV+ rRNA	RRE + rRNA
			<u>IC<sub>50</sub> (μM)</u>	<u>IC<sub>50</sub> (μM)</u>
5	81	max.	120.1	inactive
	82	max.	149.9	inactive
	83	max.	184.7	inactive
	84	max.	120.1	inactive
	85	max.	53.8	inactive
10	86	max.	200	inactive
	87	37.0	8.1	300.1
	88	80.2	11.7	937.8
	89	6.0	12.5	75.3
	90	3.0	15	44.9
15	91	46.8	17.1	800
	92	4.0	2.5	10.1
	93	max	12.7	Inactive
	94	20.8	12	249.7
	95	2.8	13.7	39
20	96	5.0	15	75
	97	22.2	17.8	394.8
	98	2.7	3	8.1
	99	max	11.5	inactive
	100	49.6	12.1	600
25	101	3.6	14	50.1
	102	3.0	15.1	45.1
	103	10.0	20	200.1
	104	15.1	5	74.9

- 50 -

	<u>Compound</u>	<u>Specificity</u>	<u>HBV+ rRNA</u> <u>IC<sub>50</sub> (μM)</u>	<u>RRE + rRNA</u> <u>IC<sub>50</sub> (μM)</u>
5	105	8.1	9.9	80.1
	106	5.8	12.1	70.1
	107	1.7	14.9	25
	108	2.5	16.1	40
	109	1.8	20	35
10	110	4.0	20	79.9
	111	20.0	23	460.3
	112	30.0	25	750.6
	113	9.1	33.1	300
	114	5.5	40	220
15	115	1.7	75.1	130.1
	116	1.3	20	25.1
	117	max.	15.6	inactive
	118	2.0	25	50.1
	119	7.1	35	250.1
20	120	24.1	46	1109.1
	121	5.0	79.9	400.1
	122	14.9	20.1	300
	123	3.0	25	74.9
	124	10.0	29.9	300
25	125	28.5	35.1	1000
	126	8.3	60.1	500.5
	127	1.0	80	80
	128	4.5	22	99.9



- 51 -

	<u>Compound</u>	<u>Specificity</u>	<u>HBV+ rRNA</u>	<u>RRE + rRNA</u>
			<u>IC<sub>50</sub> (μM)</u>	<u>IC<sub>50</sub> (μM)</u>
5	129	4.8	25	120
	130	12.2	32.9	400
	131	1.0	36.7	36.7
	132	11.4	70.1	800
	133	1.0	99.9	99.9
10	134	2.0	100.1	200
	135	2.0	149.9	300.1
	136	5.0	203.4	1017.2
	137	1.0	400	400
	138	0.1	500	49.9
15	139	0.2	2094.3	400.2
	140	4.2	120	499.9
	141	1.3	165.4	214.5
	142	5.0	226	1129.9
	143	0.5	445.9	223
20	144	0.1	1000	99.9
	145	3.1	129.9	400
	146	6.1	196	1194.9
	147	1.0	299.9	299.9
	148	max.	496.9	inactive
25	149	0.4	1000.1	399.9
	150	4.7	149.9	700.8
	151	3.5	200.0	699.9
	152	0.5	299.9	150

- 52 -

	<u>Compound</u>	<u>Specificity</u>	HBV+ rRNA <u>IC<sub>50</sub> (μM)</u>	RRE + rRNA <u>IC<sub>50</sub> (μM)</u>
5	153	1.0	499.9	499.9
	154	0.3	1180.6	300
	155	max.	25.5	inactive
	156	max.	22.8	inactive
	157	max.	2.9	inactive
10	158	max.	7.4	inactive
	159	max.	2.7	inactive
	160	44.2	9.5	420.2
	161	max.	36.1	inactive
	162	max.	11.7	inactive
15	163	max.	99	inactive
	164	max.	48	inactive
	165	max.	18.8	inactive
	166	max.	36.3	inactive
	167	max.	50.9	inactive
20	168	max.	19.7	inactive
	169	max.	15.4	inactive
	170	119.3	6.2	739.9
	171	max.	9.4	inactive
	172	max	69.2	inactive
25	173	max.	89	inactive
	174	201.1	5.6	1126.3
	175	max.	5.9	inactive

- 53 -

	<u>Compound</u>	<u>Specificity</u>	HBV+ rRNA	RRE + rRNA
			<u>IC<sub>50</sub> (μM)</u>	<u>IC<sub>50</sub> (μM)</u>
5	176	166.5	6.8	1132.5
	177	max.	6.2	inactive
	178	max.	12.7	inactive
	179	max.	13.2	inactive
	180	max.	13.7	inactive
10	181	max.	60	inactive
	182	max.	70.4	inactive
	183	max.	47.8	inactive
	184	2.0	1.7	3.4
	185	max.	203.3	inactive
15	186	max.	88.4	inactive
	187	max.	61	inactive
	188	max.	54.9	inactive
	189		inactive	inactive
	190	max.	223.7	inactive
20	191	max.	87	inactive
	192	max.	235	inactive
	193	max.	28	inactive
	194	max.	34	inactive
	195	max.	28.2	inactive
25	196	max.	32.7	inactive
	197		inactive	inactive
	198	max.	14.8	inactive

- 54 -

	<u>Compound</u>	<u>Specificity</u>	<u>HBV + rRNA</u> <u>IC<sub>50</sub> (μM)</u>	<u>RRE + rRNA</u> <u>IC<sub>50</sub> (μM)</u>
5	199	max.	33.8	inactive
	200	max.	12.1	inactive
	201	max.	53	inactive
	202	max.	68	inactive
	203	85.6	15.6	1334.6
10	204	max.	18.4	inactive
	205	max.	19.1	inactive
	206	max.	12.4	inactive
	207	max.	35.9	inactive
	208	max.	38.2	inactive
15	209	max.	9.3	inactive
	210	max.	3.3	inactive
	211	20.2	5.5	111
	212	max.	23.1	inactive
	213	33.3	15.6	518.8
20	214	max.	42.6	inactive
	215	max.	50	inactive
	216	max.	9	inactive
	217	max.	8.1	inactive
	218	12.0	4.8	57.8
25	219		inactive	inactive
	220	33.3	27.6	918.5
	221	max.	402.8	inactive

- 55 -

5

10

15

20

25

<u>Compound</u>	<u>Specificity</u>	<u>HBV + rRNA</u> <u>IC<sub>50</sub> (μM)</u>	<u>RRE + rRNA</u> <u>IC<sub>50</sub> (μM)</u>
222	max.	48	inactive
223	max.	48.8	inactive
224	max.	60.1	inactive
225	max.	193.3	inactive
226	5.0	84.4	422
227	max.	12.8	inactive
228	max.	79.3	inactive
229	max.	4.4	inactive
230	max.	20.9	inactive
231	max.	195.4	inactive
232	max.	45.5	inactive
233	max.	22.9	inactive
234	max.	48.2	inactive
235	max.	9.9	inactive
236		inactive	inactive
237	max.	51.6	inactive
238	max.	1071	inactive
239	max.	22.8	inactive
240		inactive	inactive
241	max.	5.7	inactive
242	2.5	54.2	135.6
243	max.	86.7	inactive
244		inactive	inactive

- 56 -

<u>Compound</u>	<u>Specificity</u>	<u>HBV+ rRNA</u> <u>IC<sub>50</sub> (μM)</u>	<u>RRE + rRNA</u> <u>IC<sub>50</sub> (μM)</u>
245	max.	8.1	inactive
246	max.	5.6	inactive
247	max.	6.7	inactive
248		inactive	inactive
249	max.	47.2	inactive
250		inactive	inactive
251	max.	45	inactive
252		inactive	inactive
253	59.9	11	658.8
254	3.7	87.7	328.7

**Example 2: DETERMINATION OF ANTIVIRAL ACTIVITY**

**i. Materials and Methods**

The antiviral activity of the compounds of the present invention is measured by the methods described in Korba and Gering, Antiviral Res. 19: 55-70 (1992).

Whenever possible, all materials were prepared sterile and solutions passed through 0.22 micron filters.

- 57 -

5        Materials

1.        HBV hybridization probe: A 3.2kb full genome length HBV fragment was retrieved from a restriction digest of a plasmid clone (*e.g.*, pAM6) electrophoresed in a 1% agarose gel, isolated, and stored at -20 °C.
  
- 10       2.        HBV gel standard: 1.0µl HBV standards (100 ng/ml) plus 5µl tracking dye and 14 µl TE (per lane). HBV standards were made by performing separate *Bam HI* and *Eco RI* digests of cloned HBV DNA. The digests were combined in equimolar amounts and stored at -20 °C. This produced several HBV fragment (positive hybridization controls) as well as non-HBV  
15       plasmid DNA fragments (negative hybridization control). For example, pAM6 produces HBV DNA fragments of 3.2, 1.85, and 1.35kb, as well as a plasmid fragment of 4.3kb. This mixture served as a positive and a negative hybridization control, a size standard, and a quantitation standard that occupied only one lane of a gel.
  
- 20       3.        HBV media standards: Culture medium (RP2) was collected from confluent non-G418 treated 2.2.15 cells and stored frozen at -70 °C. Aliquots were pooled as necessary (typically 1-2 liters) and centrifuged at 7000xg, 10  
25       minutes to remove cellular debris. The supernatant was added to an Amicon Inc. "stirred cell" (8400 series) fitted with a YM100 membrane (cat. No. 13642) 400 ml, and ultrafiltered at approximately 20psi nitrogen with

- 58 -

5 constant stirring, at 4°C (approximately 3-5 hours), to insure that the  
membrane was not allowed to dry. This resulted in approximately a 40-fold  
concentration by volume. HBV DNA content in the concentrated sample  
was quantitated by blot hybridization against a known standard quantity of  
HBV DNA. Approximately a 2-fold loss of HBV DNA (as compared to the  
10 theoretical starting concentration) frequently occurred during this process.  
The final product was diluted to a standard concentration of 10ng HBV  
DNA/ml with RPMI 1640 (without FBS). The concentration was then  
rechecked by blot hybridization. The adjusted standard pool was again  
rechecked by blot hybridization. The adjusted standard pool was stored at -  
15 70 °C in 5ml aliquots in screw-capped tubes and stable for at least 5 years.

ii. Guidelines for the Culture of 2.2.15 Cells

Cell cultures were handled aseptically, without antibiotics, and in  
contained facilities (BS level II). The basal culture medium used for the culture of  
20 2.2.15 cells was RPMI 1640. Fetal Bovine Serum (FBS) was added at either 2% or  
4% final concentration, and was not heat inactivated or lot tested. L-Glutamine was  
added to a final concentration of 4 mM. Complete culture medium was stored at 4  
°C for up to 4 weeks. Cells were grown and maintained at 37 °C in a 5% CO<sub>2</sub>  
humidified atmosphere.

25 Flasks and plates were routinely seeded at a density of  $3-5 \times 10^4$   
cells/cm<sup>2</sup>. This seeding density produced confluent cultures in 3-5 days. Seeding



- 59 -

5 densities of approximately  $1 \times 10^4$  cells/cm<sup>2</sup> were permissible and prolonged time to confluence. At confluence, cells were at a density of  $3-5 \times 10^5$  cells/cm<sup>2</sup> and produced their maximal and relatively stable levels of HBV. A confluent, "healthy" T-75 flask, grown in medium with 4% FBS was subcultured in up to 6, 96-well or 24-well flat bottom plates.

10

iii. "Primary" or "Screening" Assay (96-well plate format)

This assay format is well suited to the screening of test compounds for potential antiviral activity. The assay provides a threshold assessment of antiviral activity by measuring (i) the levels of HBV virion release from the cells and (ii) cytotoxicity. Two rows of cells were used for each compound, and 4 rows for the assay controls (two for untreated, and two for positive antiviral control, *e.g.*, 3TC). After incubating in the presence of test compound for 9 days, the media were harvested, transferred to 96-well U-bottom plates, and centrifuged. The supernatants were transferred to tubes for dot blot hybridization analysis of HBV virion DNA. The medium was aspirated off of the toxicity plates and discarded. Toxicity plates were then incubated with neutral red dye (MTT can also be used if preferred), washed with DPBS, developed with an acetic acid-ethanol solution, and assayed in a plate reader.

20

There are many options as to what concentration ranges to use for these assays. The compounds of the present invention were tested for toxicity at as high a concentration as the compound's solubility and the toxicity of the diluent

25

- 60 -

5 (frequently DMSO) allowed. The 2.2.15 cells will tolerate 2-3% DMSO for up to 10 days with little loss of viability.

1. 2.2.15 cells were seeded into 96-well, flat bottom tissue culture plates as described above. Duplicate plates were used for the antiviral treatments for each test compound (up to 8 compounds per pair of plates); after three to four days, cells were confluent and medium was yellow in color; the medium was removed and replaced with 100 $\mu$ l RP2 24 hours before the beginning of drug treatment.

10
- 2a. Compounds were prepared for antiviral treatment as follows:
  - i) For each compound, a total of 4 concentrations were examined; this required 9 sets of 4 sterile, 1.1 ml minitubes (36 tubes per compound). Into the first tube of each set, sufficient compound was aliquoted to make up 700  $\mu$ l (for 10-fold dilution series) or 980 $\mu$ l (for 3.3-fold dilution series) of the highest test concentration; the other tubes were left empty.

15
  - ii) The sample aliquots for the last day of treatment were set up at a 3 fold higher concentration relative to the other aliquots; this was done to provide sufficient material for DNA analysis (see below).

20
  - iii) The tubes were covered with the rack lids, labelled appropriately, and stored at -20 °C or the appropriate temperature for the test compounds; this procedure prevents

25

- 61 -

5 multiple freeze-thaw cycles of stock solutions of the test compound, and ensures that all 9 daily test aliquots are treated in an identical manner with respect to temperature variations. If compounds were stored at 4 °C, the tops of the tubes were covered with a sheet of parafilm to prevent evaporation.

10 2b. Compounds were prepared for toxicity treatment as follows:  
For each compound, a total of 4 concentrations were examined. This required 9 sets of 4 sterile, 1.1 ml minitubes (36 tubes per compound). Into the first tube of each set, sufficient compound was aliquoted to make up 465 $\mu$ l of the highest test concentration. The  
15 other tubes were left empty. The tubes were covered with the rack lids, and stored at -20 °C or at an appropriate temperature. Tubes for the last day of treatment for toxicity testing contain the same amount of compound as used the previous days (not 3x as for the antiviral assays).

20 3a. The compound dilution series was prepared for antiviral treatment as follows:

i) For a 10-fold dilution series: to the first (compound containing) tube was added 700 $\mu$ l (175 $\mu$ l x 4). 630 $\mu$ l (210 $\mu$ l x 3) RP2 culture medium was added to the remaining 3 tubes.  
25 The first tube was mixed by pipeting up and down with a pipetman (a multichannel pipetman permits the simultaneous

- 62 -

5 processing of multiple compounds). 70μl of test compound-containing medium were serially transferred from the first tube to the other 3 tubes, taking care to thoroughly mix each tube before transferring medium;

10 ii) For a 3.3-fold dilution series:  $960\mu\text{l}$  ( $160\mu\text{l} \times 6$ ) of RP2 was added to the first tube (containing the aliquot of compound) and  $640\mu\text{l}$  ( $160\mu\text{l} \times 4$ ) RP2 was added to each of the empty tubes.  $280\mu\text{l}$  of test compound-containing medium were serially transferred from the first tube to the other 3 tubes.

15 3b. The cytotoxicity of test compounds was analyzed in a 3.3-fold dilution series as follows. 485 $\mu$ l (155 $\mu$ l x 3) of RP2 were added to the first tube (containing the aliquot of compound) and 310 $\mu$ L (155 $\mu$ l x 2) RP2 to each of the 3 remaining tubes. 150 $\mu$ l of test compound-containing medium were serially transferred from the first tube to the other 3 tubes.

20

4. Treatments were initiated by the following procedure:

25

- a) The culture medium was removed with care to minimize the time that the cell monolayers are without medium;
- b) 100 $\mu$ l of each dilution of every compound was added to each of 6 wells (3 wells per plate) using the configuration listed below as an example. The lowest concentration was added

- 63 -

5 first, and the same tips were used to add the higher concentrations. The untreated cells received 100 $\mu$ l RP2 per well (untreated cells have to be carried on only one pair of the antiviral assay plates).

10	<u>Col. 1 -3</u>	<u>Col.4-6</u>	<u>Col.7-9</u>	<u>Col. 10-12</u>
	Row A----- untreated cells -----			
	Row B drug 1 @ 1X	drug 1 @ 1/10X	drug 1 @ 1/100X	drug 1 @
				/1000X
	Row C drug 2 @ 1X	.	.	.
15	Row D .	.	.	.
	Row E .	.	.	.
	Row F .	.	.	.
	Row G .	.	.	.
	Row H .	.	.	drug 7 @1/1000

20

25

- c) Treatments were repeated daily for 9 days. For the last day of treatment, and additional 200 $\mu$ l RP2 were added to each well of the antiviral assay plates after the wells are treated with the test compounds (a total of 300 $\mu$ l medium per well).

- 64 -

- 5           5.     A single plate was used for the toxicity treatments for each test  
compound (up to 7 compounds per plates since the top row were  
reserved for untreated cells on every toxicity plate). To initiate the  
treatments, the culture medium was removed, and 100 $\mu$ l of each  
dilution of every compound were added to each of 3 wells using a  
10           configuration similar to that shown above for the antiviral treatments.  
The untreated cells received 100 $\mu$ l RP2 per well. Treatment was  
repeated daily for 9 days.
- 15           6.     The assay was terminated and samples harvested for quantitative  
analysis of HBV virion DNA, by removing the culture medium 24  
hours following the 9th day of treatment and storing the culture  
medium in 96-well U-bottom culture plates. These samples were  
stored at 4 °C until blotting was performed. Samples were  
eventually be transferred to -20 °C for long term storage.

20           iv. Assay for Effects on Intracellular HBV Replication (24-well plate format)

This assay format serves to further define the action of potential  
antiviral agents by permitting an assessment of the levels of intracellular HBV DNA  
replicative forms. This type of assay is usually performed on compounds identified  
as active in the 96-well plate format since the effective antiviral concentrations  
25           observed in those experiments can be used as a guide for this type of assay (which  
is considerably more labor intensive and costly). In general, a 3- to 5-fold higher

- 65 -

5 concentration of compound will be needed than that observed in the antiviral assay to produce similar levels of effects on intracellular HBV DNA replication. 2.2.15 cells are seeded in 24-well plates for this assay and treated with test compounds for 9 days. The medium is collected at the end of the treatment period for analysis of HBV virion DNA. For analysis of intracellular HBV DNA, the monolayers are  
10 lysed with guanidine thiocyanate/sarcosyl/ $\beta$ ME, dialyzed, digested with SDS/Proteinase K, extracted with phenol and chloroform, and precipitated with sodium acetate/isopropanol. The intracellular DNAs are then resuspended, digested with *Hind* III, subjected to gel electrophoresis, and transferred to nitrocellulose for hybridization analysis.

15 1. 24-well culture plates are seeded as described above. The day before the addition of compounds, the medium is changed to RP2 (0.5 ml per well). As in the 96-well plate assay, duplicate plates are used. A total of 2 wells on each plate are treated with each dilution of compound (4 wells per dilution).

20 2. Compounds are prepared for antiviral treatment as follows:  
For each compound, a total of 4 concentrations are examined. This requires 9 sets of 4 sterile, 1.1 ml minitubes (36 tubes per compound). Into the first tube of each set, sufficient compound is aliquoted to make up 2.2ml of drug-containing medium (additional  
25 medium to make up the proper total volume was added at the time of cell treatment (see step 4 below). Tubes for the last day of treatment

- 66 -

- 5 for toxicity testing contain the same amount of compound as used the previous days (not 3X as for the 96-well plate assay). The tubes are covered with the rack lids, and stored at -20 °C or at an appropriate temperature.
- 10 3. Compounds in this assay are tested in a 3.3-fold dilution series. To make this dilution series, 720 $\mu$ l (240 $\mu$ l x 3) are added to the first tube. 460 $\mu$ l (230 $\mu$ l x 2) of medium are added to the remaining 3 tubes. 200 $\mu$ l are serially transferred from the first tube to the remaining 3 tubes.
- 15 4. Wells are treated from the lowest concentration of drug to the highest, adding 100 $\mu$ l to each of the 4 wells. An additional 400 $\mu$ l of RP2 (without drugs) are added to each well. Culture medium is continually changed and test compounds are added each day for a total of 9 days.
- 20 5. 24-hours following the final addition of compound, the culture medium is collected and stored in new 24-well plates. A 250 $\mu$ l aliquot of each stored culture medium sample is transferred to a 96-well U-bottomed culture plates (one plate can hold samples from up to 4, 24-well plates), and stored at 4 °C until dot blotting is performed.
- 25 6. The cell monolayers are then lysed for analysis of intracellular HBV DNA as described above.



- 67 -

5       vi.    Neutral Red Dye Determination of Drug Toxicity

          The antiviral effect of any compound was evaluated relative to its toxicity. Cells were seeded and treated following the guidelines above, after which a neutral red dye uptake assay was performed to assess toxicity as described below. Other assays of cytotoxicity (*e.g.*, MTT) can be substituted for the procedure described below. Note that this procedure assesses toxicity under culture and treatment conditions which are identical to those used for the antiviral analyses, thereby permitting a determination as to whether the reduction in virus production due to a specific antiviral effect or due to a cytotoxic effect on the host cell. Since the cultures, by necessity, are at confluence, the cytotoxic effects of the test compounds will probably be reduced relative to the cytotoxic effects that would be expected for actively dividing cells.

1.       Cultures were treated with test compounds on the designated toxicity plates as described above.
2.       The culture medium was carefully removed 24 hours following the 9th day of treatment. The monolayer was fully removed in the top left three wells (row A, col. 1-3). These wells were used as "blanks" for the plate reader.
3.       100 $\mu$ l DPBS (containing 0.01 % neutral red dye) were added to each well, including the empty wells, and Incubated in the tissue culture incubator for 30 minutes.

- 68 -

5                   4.     The dye was removed carefully and gently, using a multichannel  
pipetman since the monolayers can become fragile after incubation  
with neutral red dye. 200 $\mu$ l DPBS were added to all the wells taking  
care not to displace the monolayers. The DPBS was removed and  
100 $\mu$ l 50% EtOH/1% glacial acetic acid (in H<sub>2</sub>O) were added to each  
10                   well. The plates were mixed for 15 minutes on an orbital platform  
shaker (120-150 RPM) to allow for full extraction of the dye from  
the cells.

                  5.     Optical absorbance at 510nm was read in an ELISA type plate  
reader, using the 3 empty wells to set the background. An average  
15                   absorbance value was calculated for the 9 untreated cultures on a  
plate, and the absorbance of dye for the treated cultures on the same  
plate was expressed as a percentage of that value. This procedure  
was repeated for each individual plate.

20                   Table 2 shows the antiviral activity of a group of  $\epsilon$ RNA ligands. The  
micromolar concentration at which the production of extra-cellular virus was  
reduced by 50% (EC<sub>50</sub>) and was reduced by 90% (EC<sub>90</sub>) is reported. Because HBV  
does not cause cell death, the cytotoxic potential of the compounds can also be  
estimated in the same cultures by measuring cell death by the neutral red uptake  
25                   method, as described above. This value is expressed by 50% cell death (CC<sub>50</sub>  $\mu$ M).

- 69 -

5      **Table 2. Antiviral Activity of Selected 1,3,5-Triazine Compounds**

Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
1	> 100	> 100	14
2	33	83	> 100
3	4.8	17	> 100
4	3.4	10	3.7
5	0.411	1.2	307
6	0.3	0.7	29
7	33	89	29
8	> 100	> 100	196
9	0.474	1.8	175
10	> 100	> 100	> 300
11	3.9	15	27
12	0.759	4.3	77
13	8.4	208	> 300
14	> 100	> 100	> 100
15	0.5	4	0.6
16	15	218	304
18	6.5	32	105
19	> 100	> 100	> 300
20	4.1	12	7.1
21	5.6	18	5.6
22	> 100	> 100	47
23	40	106	145
24	18	148	75
25	1	9.4	105

- 70 -

	Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
5	27	> 100	> 100	> 300
	28	> 100	> 100	33
	29	1.1	6.3	15
	30	54	199	230
	33	> 100	> 100	50
10	34	66	181	41
	35	2.3	17	3.9
	38	0.592	2.7	125
	39	1	17	5.3
	40	30	79	136
15	41	36	122	> 300
	42	6.1	22	237
	43	0.841	3.9	87
	44	4.8	20	285
	46	> 100	> 100	> 100
20	49	1.2	11	140
	50	29	70	30
	51	15	73	130
	52	4.5	24	11
	53	1.2	82	72
25	56	> 100	> 100	63
	61	3.7	10	1.3
	62	> 100	> 100	> 100
	63	> 100	> 100	> 100
	65	> 100	> 100	17

- 71 -

	Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
5	67	> 100	> 100	> 100
	68	> 100	> 100	> 300
	74	> 100	> 100	> 100
	85	> 100	> 100	209
	88	4.8	14	> 100
10	91	> 100	> 100	> 300
	93	0.006	0.503	54
	94	> 100	> 100	> 300
	97	> 100	> 100	49
	99	52	162	157
15	100	39	172	57
	113	> 100	> 100	> 300
	117	4.2	36	> 300
	120	> 100	> 100	45
	125	3.9	11	265
20	155	0.001	0.063	59
	156	4	13	359
	157	0.71	2.9	59
	158	42	129	112
	159	0.498	1.9	101
25	160	0.42	4.1	> 300
	161	0.55	6.8	> 300
	162	0.923	7.5	436
	163	0.958	7.6	150
	164	0.863	9.7	201

- 72 -

	Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
5	165	3.8	11	> 300
	166	1.4	11	252
	167	1.7	12	161
	168	0.7	15	> 300
	169	2.5	37	> 300
10	170	4	40	499
	171	5	52	174
	172	5.2	64	> 300
	173	5	68	247
	174	5.5	72	133
15	175	5	75	> 300
	176	31	86	> 300
	177	5	90	> 300
	178	5	95	272
	179	26	123	> 300
20	180	12	131	> 300
	181	6	136	> 300
	182	54	191	> 300
	183	52	231	> 300
	184	13	111	441
25	185	12	107	> 300
	186	66	332	771
	187	7.1	116	266
	188	12	102	218
	189	2.5	60	88

- 73 -

	Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
5	190	> 100	> 100	945
	191	> 100	> 100	836
	192	> 100	> 100	739
	193	> 100	> 100	666
	194	> 100	> 100	626
10	195	> 100	> 100	400
	196	> 100	> 100	381
	197	> 100	> 100	354
	198	> 100	> 100	336
	199	> 100	> 100	331
15	200	> 100	> 100	309
	201	> 100	> 100	> 300
	202	> 100	> 100	> 300
	203	> 100	> 100	> 300
	204	> 100	> 100	> 300
20	205	> 100	> 100	> 300
	206	> 100	> 100	> 300
	207	> 100	> 100	> 300
	208	> 100	> 100	> 300
	209	> 100	> 100	> 300
25	210	> 100	> 100	> 300
	211	> 100	> 100	> 300
	212	> 100	> 100	> 300
	213	> 100	> 100	> 300
	214	> 100	> 100	> 300

- 74 -

	Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
5	215	> 100	> 100	> 300
	216	> 100	> 100	> 300
	217	> 100	> 100	> 300
	218	> 100	> 100	> 300
	219	> 100	> 100	> 300
10	220	> 100	> 100	> 300
	221	> 100	> 100	> 300
	222	> 100	> 100	> 300
	223	> 100	> 100	> 300
	224	> 100	> 100	> 300
15	225	> 100	> 100	281
	226	> 100	> 100	280
	227	> 100	> 100	247
	228	> 100	> 100	243
	229	> 100	> 100	237
20	230	> 100	> 100	231
	231	> 100	> 100	218
	232	> 100	> 100	190
	233	> 100	> 100	188
	234	> 100	> 100	188
25	235	> 100	> 100	178
	236	> 100	> 100	171
	237	> 100	> 100	163
	238	> 100	> 100	163
	239	> 100	> 100	154



- 75 -

Structure	<u>Antiviral</u> <u>EC<sub>50</sub> (μM)</u>	<u>Antiviral</u> <u>EC<sub>90</sub> (μM)</u>	<u>50% Cell Death</u> <u>CC<sub>50</sub> (μM)</u>
240	> 100	> 100	154
241	39	111	164
242	40	117	172
243	67	280	409
244	34	103	143
245	> 100	> 100	137
246	5.2	103	141
247	> 100	> 100	136
248	> 100	> 100	135
249	> 100	> 100	134
250	> 100	> 100	123
251	> 100	> 100	96
252	9.5	85	61
253	18	286	193
254	> 100	> 100	58

The compounds show a potent antiviral activity. In particular, compounds 5 and 6 show antiviral activities with EC<sub>90</sub> values within 3 to 5-fold of that observed for 3TC (2'-deoxy-3-thiacytidine), a well known antiviral drug, in the same assay. The figure ">" represents that no effect was observed at the highest concentration tested.

It is noted that the compounds tested above evidenced a wide range of antiviral activity, with some compounds demonstrating more activity than other

- 76 -

5 compounds. It is believed that all structures 1-254 may have activity against the HBV. However, as is understood by persons of ordinary skill in the art, assay results demonstrating inactivity may be due to a number of factors, such as cell permeability and metabolic stability.

10 **Example 3: Antiviral Combination Therapies**

Compound 5 and 3TC were tested at varying concentrations, alone and in combination, in the antiviral assay outlined in Example 2, above. The reduction in virus production when these drugs were tested independently is shown in Figure 3. 3TC alone causes 90% viral reduction ( $EC_{90}$ ) at 216 nM, while  
15 compound 5 alone exhibits an  $EC_{90}$  at 1300 nM.

When compounds 5 and 3TC were tested in combination with molar ratios of 1:15, 1:5, and 1:1.5, the results are shown in Figures 4A, 4B, and 4C. The front row column of each plot shows the expected % viral reduction if the effects of each drug are additive. The back row in each plot shows the observed  
20 antiviral activity. The results show a strong synergistic effect when the drugs are mixed at a 1:15 (3TC:compound 5) molar ratio. For example, approximately 7% viral reduction is the expected additive effect of 20 nM 3TC and 300 nM compound 5, but the observed viral reduction reached 92%, which is indicative of a synergistic interaction.

- 77 -

5                    These results suggest that when combined in a 1:15 ratio these two drugs could be used at approximately 4 to 10-fold lower concentrations than those needed when each drug is administered alone.

**Example 4: Nucleic Acid Binding of Triazine Compounds**

10                    Reaction mixtures (10  $\mu$ l) containing 50 mM Tris-HCl, pH 7.5, 50 mM KCl, 0.5 mg/ml rRNA (as a carrier), 0.5 pmol of 3'-end  $^{32}$ P-labeled  $\epsilon$ RNA, 0.05 units of *B. Cereus* RNase, and compound 5 at concentrations between 0 and 200  $\mu$ M were incubated at 25  $^{\circ}$ C for 30 minutes. The reaction products were resolved by gel electrophoresis in polyacrylamide gels. *B. Cereus* nuclease is a  
15                    single strand specific RNase. The results are shown in Figure 5A, wherein the sections of the ladder of digestion products corresponding to cleavage events occurring in the bulge and loop sections of the  $\epsilon$ RNA structure. It is evident that increasing concentrations (indicated at top) of compound 5 in the reaction results in a decrease in cleavage at specific positions of the bulge indicated by stars.  
20                    Conversely, the loop and other regions of the structure where not affected. Similar results were obtained using compounds 6, 15, 28, 33, and 61.

                      These results indicate that the triazine compound bind RNA and that it does so at a site overlapping the bulge region. Moreover, an RNA containing a structural element similar to the bulge of  $\epsilon$ RNA is likely to be bound by a triazine  
25                    compound increasing the stability of the RNA structure.

- 78 -

5                   Further, a smaller RNA target containing only the bulge region of  $\epsilon$ RNA was prepared, and the melting temperature ( $t_m$ ) of this RNA structure was determined by monitoring the change in optical density (OD) at 260 nm while increasing the temperature. A 1 ml aliquot of reaction mixture contained 50 mM NaCl, 10 mM sodium cacodylate, pH 7.0, 10% DMSO, and 2  $\mu$ M bulge RNA.

10                   The results are shown in Figure 6. The y-axis shows the first derivative of the change in OD at 260 nm in OD units/degree C. The reaction containing 2  $\mu$ M RNA and no drug shows that the RNA has a  $t_m$  of 65 °C, and upon addition of 10  $\mu$ M compound 6, the  $t_m$  is shifted to 83 °C. These results suggest that the RNA structure is stabilized by the ligand.


15                   All patents, applications, test methods, and publications mentioned herein are hereby incorporated by reference.

                  Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed disclosure. All such modifications are within the full extended scope of the appended claims.


- 79 -

5      What is claimed is:

1.            A pharmaceutical formulation comprising a compound of formula IA:



or IB:



10

15

20

- 80 -

5

wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, non-aromatic heterocyclic, fused or polycyclic ring and aryloxy;

10

wherein said alkyl, alkenyl or alkynyl is optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl or alkynyl, and

15

wherein said aryl, aryloxy, heteroaryl, non-aromatic heterocyclic or fused or polycyclic ring is optionally substituted by one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amides, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

20

or wherein  $R^1$  and  $R^2$  together,  $R^3$  and  $R^4$  together, or  $R^5$  and  $R^6$  together, optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring, said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester,

25

- 81 -

5       amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl,  
cycloalkenyl and alkynyl;

                  or wherein R<sup>7</sup> and R<sup>8</sup> together optionally form a cycloalkyl,  
cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring wherein said  
cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring are  
10       optionally substituted with one or more substituents selected from the group  
consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy,  
amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines,  
cyano, cycloalkyl, alkenyl and alkynyl, with the proviso that when R<sup>7</sup> and R<sup>8</sup>  
together form a fused or polycyclic ring, the moiety of the fused or polycyclic ring  
15       that binds with N is non-aromatic;

                  and pharmaceutically acceptable salts thereof;

                  and a pharmaceutically acceptable carrier or diluent.

2.       The formulation of claim 1 comprising a compound of  
20       formula IB wherein one R<sup>1</sup> or R<sup>2</sup> is an optionally substituted aryl.

3.       For formulation of claim 1 comprising a compound of  
formula IB wherein one of R<sup>7</sup> or R<sup>8</sup> is an optionally substituted aryl.

25       4.       The formulation of claim 1, said formulation further  
comprising at least one component selected from the group consisting of a diluent,

- 82 -

5       excipient, adjuvant, one or more non-triazine based antiviral or antibiotic agents,  
and mixtures thereof.


5.       The formulation of claim 1, wherein said compound binds one  
or more nucleic acids.

10

6.       The formulation of claim 5, wherein said one or more nucleic  
acids are RNA.


7.       A method of preventing or treating hepatitis B virus infection  
15       in a patient in need of such treatment, said method comprising administering a  
compound of the formula IA:

20



or IB:

25





- 83 -

5

wherein  $R^1$ ,  $R^2$ ,  $R^3$ ,  $R^4$ ,  $R^5$ ,  $R^6$ ,  $R^7$  and  $R^8$  are each independently selected from the  
10 group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, non-  
aromatic heterocyclic, fused or polycyclic ring and aryloxy;

wherein said alkyl, alkenyl or alkynyl is optionally substituted  
with one or more substituents selected from the group consisting of halogen,  
hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl,  
15 carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl,  
alkenyl, cycloalkenyl or alkynyl, and

wherein said aryl, aryloxy, heteroaryl, non-aromatic  
heterocyclic, or fused or polycyclic ring is optionally substituted by one or more  
substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro,  
20 trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide,  
primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and  
alkynyl;

or wherein  $R^1$  and  $R^2$  together,  $R^3$  and  $R^4$  together, or  $R^5$  and  $R^6$   
together, optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic,  
25 heteroaryl, or fused or polycyclic ring, said cycloalkyl, cycloalkenyl, non-aromatic  
heterocyclic, heteroaryl, or fused or polycyclic ring optionally substituted with one

- 84 -

5 or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

or wherein R<sup>7</sup> and R<sup>8</sup> together optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring wherein said  
10 cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring are optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl and alkynyl, with the proviso that when R<sup>7</sup> and R<sup>8</sup>  
15 together form a fused or polycyclic ring, the moiety of the fused or polycyclic ring that binds with N is non-aromatic;

and pharmaceutically acceptable salts thereof;

and a pharmaceutically acceptable carrier or diluent, wherein said  
20 compound is administered in an amount and for a time sufficient to inhibit viral replication.

8. The method of claim 7, wherein said method comprises administering a compound of formula IB wherein one of R<sup>1</sup> or R<sup>2</sup> is an optionally  
25 substituted aryl.

- 85 -


5                    9.     The method of claim 7, wherein said method comprises administering a compound of formula IB wherein one of R<sup>7</sup> or R<sup>8</sup> is an optionally substituted aryl.

10                   10.    The method of claim 7, further comprising administering at least one component selected from the group consisting of one or more additional nontriazine based antiviral agents, and mixtures thereof.

15                   11.    The method of claim 7, wherein said antiviral formulation is administered orally.

12.    The method of claim 7, wherein said compound is administered at a dosage of 0.1 to 250 mg/Kg/day.

20                   13.    A method of preventing or treating microbial infection in a patient in need of such treatment, said method comprising administering a compound of the formula IA:




- 86 -

5

or IB:

10



wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are each independently selected  
15 from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, non-  
aromatic heterocyclic, fused or polycyclic ring and aryloxy;

wherein said alkyl, alkenyl or alkynyl is optionally substituted with  
one or more substituents selected from the group consisting of halogen, hydroxy, alkyl,  
nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide,  
20 primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl or  
alkynyl, and

wherein said aryl, heteroaryl, non-aromatic heterocyclic or fused or  
polycyclic ring is optionally substituted by one or more substituents selected from the  
group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy,  
25 amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano,  
cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

- 87 -

5                   or wherein R<sup>1</sup> and R<sup>2</sup> together, R<sup>3</sup> and R<sup>4</sup> together, or R<sup>5</sup> and R<sup>6</sup> together, optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring, said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl or fused or polycyclic ring optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, 10   alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

                  or wherein R<sup>7</sup> and R<sup>8</sup> together optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring wherein said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, or fused or polycyclic ring are optionally 15   substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl and alkynyl, with the proviso that when R<sup>7</sup> and R<sup>8</sup> together form a fused or polycyclic ring, the moiety of the fused or polycyclic ring that binds with N is non-aromatic;

20                   and pharmaceutically acceptable salts thereof;

                  and a pharmaceutically acceptable carrier or diluent wherein said compound is administered in an amount and for a time sufficient to inhibit viral replication.

14.     The method of claim 13, wherein said method comprises 25   administering a compound of formula IB wherein one of R<sup>1</sup> or R<sup>2</sup> is an optionally substituted aryl.

- 88 -

5                    15.    The method of claim 13, wherein said method comprises  
administering a compound of formula IB wherein one of R<sup>7</sup> or R<sup>8</sup> is an optionally  
substituted aryl.

                  16.    The method of claim 13, wherein said compound is co-administered  
10    with at least one component selected from the group consisting of one or more additional  
non-triazine based antibiotic agents and mixtures thereof.


                  17.    The method of claim 13 wherein said compound is administered  
orally.

15


                  18.    The method of claim 13 wherein said compound is administered at a  
dosage of 0.1 to 250 mg/Kg/day.

- 89 -

- 5                    19. A method of detecting a target nucleic acid comprising
- (a) contacting the target nucleic acid with a compound of the formula IA:



or IB:



20

wherein R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup>, R<sup>7</sup> and R<sup>8</sup> are each independently selected from the group consisting of hydrogen, alkyl, alkenyl, alkynyl, aryl, heteroaryl, non-aromatic heterocyclic, fused or polycyclic ring and aryloxy;

wherein said alkyl, alkenyl or alkynyl is optionally substituted with

25 one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide,

- 90 -

5 primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl or alkynyl, and

wherein said aryl, aryloxy, heteroaryl, non-aromatic heterocyclic, or fused or polycyclic ring is optionally substituted by one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy,  
10 alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

or wherein R<sup>1</sup> and R<sup>2</sup> together, R<sup>3</sup> and R<sup>4</sup> together, or R<sup>5</sup> and R<sup>6</sup> together, optionally form a cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or fused or polycyclic ring, said cycloalkyl, cycloalkenyl, non-aromatic heterocyclic, heteroaryl, or  
15 fused or polycyclic ring optionally substituted with one or more substituents selected from the group consisting of halogen, hydroxy, alkyl, nitro, trihalomethyl, aryl, aryloxy, alkoxy, amino, carbonyl, carboxyl, ester, amide, primary, secondary or tertiary amines, cyano, cycloalkyl, alkenyl, cycloalkenyl and alkynyl;

and pharmaceutically acceptable salts thereof;

20 and a pharmaceutically acceptable carrier or diluent; and

(b) monitoring an interaction between the target nucleic acid and at least one compound.

20. The method of claim 19, wherein said compound is labeled with a  
25 moiety selected from the group consisting of a fluorescent compound, an antibody for the target nucleic acid, and a radioactive label.






FIG. 1

FIG. 2





FIG. 3

1/1/18




FIG. 4

FIG. 5




FIG. 6




FIG. 7




FIG. 8





FIG. 9



1/3/18




FIG. 10




FIG. 11




FIG. 12

1/4/18

FIG. 13





FIG. 14




FIG. 15

1/ 5/18




FIG. 16




FIG. 17




FIG. 18

1/6/18




FIG. 19




FIG. 20




FIG. 21

1/7/18




FIG. 22





FIG. 23

FIG. 24








FIG. 25




FIG. 26




FIG. 27

2/1/18




FIG. 28





FIG. 29

FIG. 30



2/2/18

FIG. 31





FIG. 32




FIG. 33

2/3/18




FIG. 34

FIG. 35





FIG. 36



2/4/18

FIG. 37




FIG. 38





FIG. 39



SUBSTITUTE SHEET (RULE 26)

2/5/18

FIG. 40




FIG. 41





FIG. 42



2/6/18




FIG. 43

FIG. 44





FIG. 45

2/7/18




FIG. 46

FIG. 47







FIG. 48






**FIG. 50**



**FIG. 51**



3/1/18




FIG. 52





FIG. 53

FIG. 54



3/2/18




FIG. 55

FIG. 56





FIG. 57

3/3/18

FIG. 58




FIG. 59





FIG. 60



3/4/18




FIG. 61

FIG. 62





FIG. 63

3/5/18




FIG. 64

FIG. 65





FIG. 66



3/6/18




FIG. 67

FIG. 68





FIG. 69

3/7/18




FIG. 70

FIG. 71





FIG. 72








FIG. 73

FIG. 74





FIG. 75



4/1/18




FIG. 76

FIG. 77





FIG. 78

4/2/18

FIG. 79




FIG. 80





FIG. 81



4/3/18

FIG. 82




FIG. 83





FIG. 84



4/4/18




FIG. 85

FIG. 86





FIG. 87

4/5/18




FIG. 88




FIG. 89




FIG. 90

4/6/18




FIG. 91

FIG. 92





FIG. 93



4/7/18

FIG. 94




FIG. 95




FIG. 96






FIG. 97




FIG. 98





FIG. 99



5/1/18

FIG. 100




FIG. 101





FIG. 102



5/2/18




FIG. 103

FIG. 104





FIG. 105



5/3/18

FIG. 106




FIG. 107





FIG. 108



5/4/18




FIG. 109

FIG. 110





FIG. 111

5/5/18

FIG. 112




FIG. 113





FIG. 114



5/6/18

FIG. 115




FIG. 116





FIG. 117



5/7/18

FIG. 118




FIG. 119




FIG. 120






FIG. 121




FIG. 122





FIG. 123



6/1/18




FIG. 124





FIG. 125

FIG. 126



6/2/18

FIG. 127




FIG. 128





FIG. 129



6/3/18

FIG. 130




FIG. 31





FIG. 132



6/4/18

FIG. 133




FIG. 134





FIG. 135



6/5/18




FIG. 136

FIG. 137





FIG. 138

6/6/18

FIG. 139




FIG. 140





FIG. 141






FIG. 142




FIG. 143




FIG. 144



FIG. 145




FIG. 146





FIG. 147



7/1/18




FIG. 148

FIG. 149





FIG. 150

7/2/18




FIG. 151





FIG. 152

FIG. 153



7/3/18

FIG. 154





FIG. 155



7/4/18

FIG. 156





FIG. 157



7/5/18




FIG. 158




FIG. 159




FIG. 160

7/6/18

FIG. 161





FIG. 162



7/7/18




FIG. 163

FIG. 164





FIG. 165



7/8/18

FIG. 166




FIG. 167





FIG. 168






FIG. 169




FIG. 170




FIG. 171

8/1/18




FIG. 172




FIG. 173




FIG. 174




FIG. 175




FIG. 176




FIG. 177

8/3/18

FIG. 178




FIG. 179





FIG. 180



SUBSTITUTE SHEET (RULE 26)

8/4/18

FIG. 181




FIG. 182





FIG. 183



8/5/18




FIG. 184




FIG. 185




FIG. 186

**8/6/18**




FIG. 187

FIG. 188





FIG. 189





8/7/18

FIG. 190




FIG. 191





FIG. 192



9/18

FIG. 193




FIG. 194




FIG. 195





FIG. 196



9/1/18




FIG. 197




FIG. 198




FIG. 199

9/2/18

FIG. 200




FIG. 201





FIG. 202



9/3/18




FIG. 203




FIG. 204




FIG. 205

9/4/18




FIG. 206




FIG. 207




FIG. 208




FIG. 209

FIG. 210





FIG. 211



9/6/18

FIG. 212




FIG. 213





FIG. 214





10/18

FIG. 215




FIG. 216





FIG. 217



10/1/18




FIG. 218

FIG. 219





FIG. 220

10/2/18

FIG. 221




FIG. 222





FIG. 223



10/3/18

FIG. 224




FIG. 225





FIG. 226



10/4/18

FIG. 227




FIG. 228




FIG. 229




FIG. 230




FIG. 231





FIG. 232



10/6/18

FIG. 233




FIG. 234





FIG. 235



10/7/18




FIG. 236

FIG. 237





FIG. 238





11/18

FIG. 239




FIG. 240





FIG. 241



11/1/18




FIG. 242

FIG. 243





FIG. 244



11/2/18

FIG. 245




FIG. 246





FIG. 247



11/3/18




FIG. 248




FIG. 249




FIG. 250

11/4/18




FIG. 251





FIG. 252

FIG. 253



SUBSTITUTE SHEET (RULE 26)

12/18

FIG. 254




FIG. 255A




FIG. 255B





FIG. 255C



14/18

FIG. 256







FIG. 257A



15/1/18

FIG. 257B






FIG. 257C



17/18

FIG. 258 A



17/1/18






FIG. 258B

FIG. 259A



18/1/18

FIG. 259B



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/00945

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D251/66 C07D251/70 A61K31/53 C07D403/12 C07D405/12  
C07D409/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 3 549 760 A (ALBERT JAMES R ET AL) 22 December 1970 see claim 1 ---	1
X	EP 0 240 854 A (HOECHST AG) 14 October 1987 see claim 1; examples ---	1
X	EP 0 092 479 A (ADIR) 26 October 1983 see claim 1; examples ---	1
X	WO 93 20056 A (JARMAN MICHAEL ; COLEY HELEN MARY (GB)) 14 October 1993 see claim 1; examples ---	1
	--- -/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

22 April 1999

Date of mailing of the international search report

06/05/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/00945

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 505 220 A (JARMAN MICHAEL ;WILMAN DERRY EDWARD VINCENT (GB); JUDSON IAN ROBER) 23 September 1992 see claim 1; examples ---	1
X	DE 29 30 039 A (AMERICAN CYANAMID CO) 21 February 1980 see claim 1 ---	1
X	EP 0 629 622 A (ZENYAKU KOGYO KK) 21 December 1994 see claim 1; examples ---	1
X	WO 93 10116 A (PFIZER) 27 May 1993 see claim 1; examples ---	1
X	H. BRUHIN ET AL: "Antituberculosis activity of some nitrofurran derivatives" J. PHARM. PHARMACOL., vol. 21, no. 7, 1969, pages 423-433, XP002100043 see tables ---	1
X	CHEMICAL ABSTRACTS, vol. 126, no. 12, 24 March 1997 Columbus, Ohio, US; abstract no. 152444, COLEY, HELEN M.: "N-(Hydroxymethyl) melamines" XP002100046 see abstract & GEN. PHARMACOL. (1997), 28(2), 177-182 ;ISSN: 0306-3623,1997, ---	1
X	CHEMICAL ABSTRACTS, vol. 100, no. 9, 27 February 1984 Columbus, Ohio, US; abstract no. 61386, SU, YONGZHUANG ET AL: "Structure-activity relations of antitumor hexamethylmelamine derivatives" XP002100047 see abstract & ZHONGGUO YAOLI XUEBAO (1983), 4(4), 276-80 ,1983, ---	1
	---	

-/--

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/00945

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 88, no. 17, 24 April 1978 Columbus, Ohio, US; abstract no. 115093, RUTTY, C. J. ET AL: "In vitro studies with hexamethylmelamine" XP002100048 see abstract & BIOCHEM. PHARMACOL. (1977), 26(24), 2385-91 ; ISSN: 0006-2952, 1977, ---	1
X	MATSUNO, TOSHIYUKI ET AL: "Synthesis and aromatase-inhibitory activity of imidazolyl-1,3,5-triazine derivatives" CHEM. PHARM. BULL. (1997), 45(2), 291-296 CODEN: CPBTAL; ISSN: 0009-2363, 1997, XP002100044 see table 1 ---	1
X	JARMAN, MICHAEL ET AL: "Synthesis and cytotoxicity of potential tumor-inhibitory analogs of trimelamol (2,4,6-tris(hydroxymethyl)methylamino-1, 3,5-triazine) having electron-withdrawing groups in place of methyl" J. MED. CHEM. (1993), 36(26), 4195-200 CODEN: JMCMAR; ISSN: 0022-2623, 1993, XP002100045 see page 4195 - page 4196 ---	1
A	US 5 120 843 A (AYER DONALD E ET AL) 9 June 1992 see column 32, line 12; example 1 ---	1,7,13
A	EP 0 775 487 A (NIPPON SHINYAKU CO LTD) 28 May 1997 -----	1,7,13

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/00945

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 7-18  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 7-18  
are directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☒ Claims Nos.: not applicable  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

Claims Nos.: not applicable

In view of the extremely broad Markush claim 1, the search was executed with due regard to the PCT Search Guidelines (PCT/GL/2), C-III, paragraph 2.1, 2.3 read in conjunction with 3.7 and Rule 33.3 PCT, i.e. particular emphasis was put on the inventive concept, as illustrated by the examples. The international search was, in so far as possible and reasonable, complete in that it covered the entire subject-matter to which the claims are directed.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/00945

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 3549760	A	22-12-1970	NONE	
EP 0240854	A	14-10-1987	DE 3611427 A AU 7105687 A DK 171587 A FI 871448 A JP 62240673 A PH 22967 A PT 84618 A,B	08-10-1987 08-10-1987 06-10-1987 06-10-1987 21-10-1987 10-02-1989 01-05-1987
EP 0092479	A	26-10-1983	FR 2525597 A AT 17724 T AU 1381483 A CA 1203803 A JP 1432674 C JP 58189168 A JP 62040357 B OA 7400 A PT 76575 A,B US 4514399 A ZA 8302778 A	28-10-1983 15-02-1986 27-10-1983 29-04-1986 24-03-1988 04-11-1983 27-08-1987 30-11-1984 01-05-1983 30-04-1985 28-12-1983
WO 9320056	A	14-10-1993	AT 168105 T AU 676677 B AU 3894293 A DE 69319590 D DE 69319590 T EP 0632805 A ES 2118945 T JP 7505380 T US 5534625 A US 5854244 A	15-07-1998 20-03-1997 08-11-1993 13-08-1998 12-11-1998 11-01-1995 01-10-1998 15-06-1995 09-07-1996 29-12-1998
EP 0505220	A	23-09-1992	AU 1566792 A CA 2106868 A WO 9216513 A JP 6511229 T MX 9201258 A	21-10-1992 23-09-1992 01-10-1992 15-12-1994 01-10-1992
DE 2930039	A	21-02-1980	AU 528798 B AU 4738879 A BE 877950 A CH 641959 A PH 18639 A ZA 7902562 A	12-05-1983 07-02-1980 28-01-1980 30-03-1984 23-08-1985 27-08-1980
EP 0629622	A	21-12-1994	AU 3575193 A DE 69322076 D US 5489591 A CA 2131004 A WO 9317009 A	13-09-1993 17-12-1998 06-02-1996 02-09-1993 02-09-1993
WO 9310116	A	27-05-1993	AU 2599992 A EP 0613477 A FI 942188 A HU 69722 A JP 7506084 T MX 9206468 A	15-06-1993 07-09-1994 11-05-1994 28-09-1995 06-07-1995 01-05-1993

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/00945

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9310116 A		NO 941785 A PT 101046 A US 5536722 A ZA 9208683 A	11-05-1994 28-02-1994 16-07-1996 11-05-1994
US 5120843 A	09-06-1992	AU 624788 B AU 1709888 A CA 1338012 A DK 533589 A EP 0293078 A EP 0358676 A EP 0487510 A JP 7103118 B JP 2503198 T KR 9601920 B WO 8808424 A	25-06-1992 02-12-1988 30-01-1996 26-10-1989 30-11-1988 21-03-1990 27-05-1992 08-11-1995 04-10-1990 06-02-1996 03-11-1988
EP 0775487 A	28-05-1997	AU 703263 B AU 3192095 A BR 9508539 A CA 2197091 A HU 77735 A WO 9604914 A	25-03-1999 07-03-1996 28-10-1997 22-02-1996 28-07-1998 22-02-1996